

IMPORTANT: READ THE LABEL INSTRUCTIONS AND DIRECTIONS FOR USE BEFORE OPENING AND USING. ALBAUGH AUSTRALIA PTY LTD Level 1, 530 Little Collins Street, Melbourne VIC 3000 Tel 03 9909 7183 www.albaugh.com/au

C-270-041-J

used. Rates above refer to sprayed hectares and not field hectares. for further Radish, Wild Turnip, Yellow Wood, Sorrel. Some tree damage may occur, especially with post-plant application. Recovery is normally rapid. Directed application is preferred for information on Weed suppression at higher rate: Soursob, Wild Oats post-planting application especially when the higher rates are used. tree species] Weeds suppressed WA: Annual Ryegrass, Brome Grass, Dock, Doublegee, Radish, Sorrel, Soursob, Wild Turnip, Wireweed As above and including

All States

5kg or 500g/ 100L

5kg or 500g/ 100L

Provides seasonal control of annual weed germinations. Apply to bare moist ground. Non-crop uses, Commercial, Industrial, Rights of Way, Barnyard Grass, Summer Grass 10kg or 1kg/100L 24kg or 2.4kg/100L Provides long residual control in Winter rainfall areas. Apply to bare moist ground. 24kg or 2.4kg/100L 48kg or 4.8kg/100L Provides long residual control in Summer rainfall areas. Public Utility areas, Road shoulders, Drains, Headlands, Driveways, Railway tracks, Apply to bare moist ground. Read remarks under Crop Safety. Aerodromes, Gutters, Footpaths Filamentous Blue- Green Algae WA only 2g per 100 litres of water Mix in a convenient amount of water and apply when algae development first noticed. Dams, Tanks, Troughs NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION WITHHOLDING PERIODS All crops: NOT REQUIRED WHEN USED AS DIRECTED DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION b Clover: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION. DO NOT CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION kpeas: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION Faba Beans: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION repeatedly. These resistant weeds will not be controlled by Albaugh ROMAZINA 900 WG Herbicide or other Group 5 herbicides. Albaugh ROMAZINA 900 WG Herbicide is a pre-emergent herbicide, which is most effective when applied to bare, moist soil.

Since the occurrence of resistant weeds is difficult to detect prior to use, Albaugh Australia Pty Ltd accepts no liability for any losses Because it acts mainly through the roots of weeds, best results are obtained when rain or irrigation move the herbicide down to that may result from the failure of Albaugh ROMAZINA 900 WG Herbicide to control resistant weeds. the root zone within 2 weeks of application. The duration and effectiveness of control depends on the amount applied, the type of PRECAUTIONS Integrated Weed Management Strategy for TT-Canola Integrated Weed Management Strategy for TT-Canola (the Strategy) has been developed and funded by the Canola

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS sociation of Australia in co-operation with certain companies. The Strategy is available from the Canola Association of Australia. DO NOT spray foliage of desirable plants, except lupins (WA only). DO NOT use under peach, apricot, plum trees, ornamental program has been developed that outlines sound agronomic practices and integrated weed management programs designed to prunus or near newly planted shrubs. DO NOT plant crops other than those recommended for at least nine (9) months following timise the performance of TT-Canola. It is advised that consultation on IWM be undertaken with an accredited agronomist prior treatments at rates up to 2.5 kg/ha. At higher rates planting may not be possible for much longer periods afterwards. DO NOT use beet, lettuce, cabbage, tomato, cucumber and carrots are particularly sensitive to Simazine residues. • Avoid dry sowing in heavily weed infested paddocks. Wait for a weed germination after the opening rains in weedy paddocks. DO NOT exceed highest rate recommended for any particular crop. DO NOT use on white or grey sands (WA only). DO NOT use Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage. in channels or drains where the roots of desirable plants may extend. DO NOT apply under weather conditions or from spraying equipment, which may cause spray to drift onto nearby susceptible plant/crops, cropping land or pastures. Broadleaf Weeds and Ryegrass: Use Albaugh ROMAZINA 900 WG Herbicide or atrazine herbicide plus trifluralin pre-emergence. Araucaria, Corymbia, Eucalyptus and Pinus Plantations: Field trials and use experience has indicated that the following species are tolerant. Test tolerance before treating other species. **Broadleaf Weeds only:** Use atrazine herbicide post-emergence. Araucaria - all commercial species. • DO NOT use atrazine or Albaugh ROMAZINA 900 WG Herbicide if the area to be treated had a triazine herbicide applied to it Corymbia - maculate. Eucalyptus – agglomerata, calophylla, camaldulensis, cladocalyx, cloeziana, dunnii, globulus, grandis, laevopinea, nitens, pilularis, • Watch for escapes especially in paddocks with a long history of Group 5 herbicide use. pseudoglobulus, regnans, rudis, saligna, wandoo. Pinus – caribaea and hybrids, elliottii, pinaster, radiata. • DO NOT use Group 5 herbicides in consecutive years. To avoid Triazine carry-over: On acid soils (pH less than 6.5) – The maximum rate of atrazine or simazine or a combination of the 2 products to be applied to the If sufficient rain to thoroughly wet the soil through the weed root zone has not fallen within 14 days of application, water in with the equivalent of 13mm of rain. If applied under dry conditions and irrigation is not available, mechanical incorporation crop during the growing season is 2.2kg ai/ha. On alkaline soils (pH greater than 6.5) - The maximum rate of atrazine or simazine or a combination of the 2 products to be to a depth of not more than 4cm within 7 days of sowing may assist results. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT applied to the crop during the growing season is 1.1kg ai/ha. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. DO NOT use the container for any Post-emergence use - It is recommended that atrazine only be used, and at rates of 1.1kg ai/ha or less, on both acid or purpose. This product is very highly toxic to algae and aquatic macrophytes. RESISTANT WEEDS REPORTING STORAGE AND DISPOSAL Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine and simazine may be Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, resistant, get them tested and seek professional advice. replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, Half fill the tank with clean water. Commence vigorous agitation. Add granules steadily. bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of DO NOT dump. Ensure complete dispersion before filling to volume with water. Continue agitation during spraying. waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not DO NOT leave spray solution in tank overnight. burn empty containers or product. se a boom sprayer or a knapsack sprayer for small areas. 50–200L of mixture per hectare should be applied with the boom, and a SAFETY DIRECTIONS Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use. minimum equivalent to 500L per hectare through the knapsack sprayer. ean spray equipment immediately after use with a proprietary spray tank and equipment cleaner. First aid is not generally required. If in doubt, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand DO NOT apply to TT-canola by aircraft. Apply only with a low boom sprayer with a 60m buffer zone downwind of treated fields to 0800 764 766 or a doctor). natural or impounded lakes or dams, and a 20m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to

SAFETY DATA SHEET

Apply only to where run-off is unlikely to occur to where run-off may be captured by farm earthworks. TERMS AND CONDITIONS OF SUPPLY, SALE AND USE s product may be mixed with amitrole, atrazine (900g/kg), diuron, glyphosate, diquat and paraquat-diquat mixtures. Because Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop ducts may vary from time to time, a small-scale compatibility test should be carried out before mixing in the spray tank. variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and ay be used. Half fill the spray tank with clean water before adding this product at the recommended rate and mix thoroughly. manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe en add 360g/L glyphosate product at 3L/ha or 700mL/100L and the remaining water. Maintain adequate agitation and use the ixture promptly. DO NOT use lower rates of glyphosate (360g/L) where control of perennial weeds is desired. application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application, or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether baugh ROMAZINA 900 WG Herbicide is a member of the Triazines group of herbicides. Simazine has the inhibitors of notosynthesis at photosystem II mode of action. For weed resistance management Simazine is a Group 5 herbicide. Some aturally occurring weed biotypes resistant to Simazine and other Group 5 herbicides may exist through normal genetic variability any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used Suspected of causing cancer. Very toxic to aquatic life. Very toxic to aquatic life with long lasting effects. ain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid releas he environment. Wear protective gloves/protective clothing/eye protection/face protection/hearing protection. IF exposed oncerned: Get medical advice/attention. Collect spillage. Store locked up. Dispose of contents/container to hazardous or cial waste collection point, in accordance with local, regional, national and/or international regulation. IN A TRANSPORT EMERGENCY POLICE OR FIRE BRIGADE

Albaugh ROMAZINA 900 WG Herbicide may be applied prior to or after planting.

spraying eucalyptus plantations on sandy, low organic matter soils in WA & SA.

On gravely, coarse textured soils use 2.8 to 4.0 kg/ha and for sandy low organic soils use 1.6 to 3.0 kg/ha.

On sandy soils allow 2 weeks and at least 50mm of rain between spraying and planting. Use a maximum rate of 1.6 kg/ha for over-

Apply to bare ground. Best results will be achieved if applied to moist, finely tilled soil. If weeds are present add a knockdown or other

partner herbicide as a tank mix. If spraying after planting check tree tolerance to knockdown or other partner herbicide before using

Application in a band 1.5 to 2.0 metres wide, over the planting row is recommended although broadcast application can also be

For short term weed control or when applied with other pre-emergent herbicides use half rates according to the situation.

Situation & Crop | Weeds Controlled

Forestry Pre-emergence control of many annual and some All States

Including farm perennial grasses and broadleaved weeds including; tree plantations Amaranthus, Annual (Wimmera) Ryegrass, Barley Grass,

inus, Araucaria, Bindii, Brome Grass, Capeweed, Chickweed, Common

forymbia and Sowthistle, Corn Gromwell (Sheep Weed), Creeping

Eucalyptus tree Oxalis, Fat Hen, Fumitory, Ivy-leaf Speedwell, Mustard, plantings pre- and Native Geranium, Nettles, Potato Weed, Powell's

post-planting Amaranth, Red Root Amaranth, Red Shank, Shepherds

Instructions Variegated, Turnips, Turnip Weed, Wild Mustard, Wild

[see General Purse, Summer Grass, Thistles - Annual, Soldier,

PBOM 11x5.5x29 .067

013 ALB Romazina 900 WG_15kg bag_250722.indd 1