Top Print

Live Copy Area

Back

ALBAUGH TEBIX 200 GR HERBICIDE

RESTRAINTS
DO NOT apply this product to salt or erosion prone areas.
DO NOT apply this product within 100 metres of a recognised water course.
DO NOT apply Albaugh TEBIX 200 GR Herbicide on land with a slope greater than 20 percent (11 degrees).
DO NOT apply Albaugh TEBIX 200 GR Herbicide under conditions which will cause pellet movement to non-target areas during application.
RETAIN at least 20 percent of the original tree population in wildlife corridors that are at least 100 metres wide. These are essential for shade and protection of livestock and native fauna. Ideally, shelter belts should traverse variable terrain, particularly ridgelines and hill tops as well as water courses, and link other vegetated corridors where possible.
FOR LONG TERM VIABILITY retain a minimum of 5 hectares for shade areas, regardless of vegetation type.
DO NOT apply Albaugh TEBIX 200 GR Herbicide on field crops, near desirable trees or shrubs or to areas into which their roots may extend or in locations where the chemical may be washed into contact with their roots as injury or death may occur.
DO NOT apply Albaugh TEBIX 200 GR Herbicide to land to be put under field crops within 5 years of application.
SPRAY DRIFT RESTRAINTS
Specific definitions for terms used in this section of the label can be found at apvma.gov.au/ spraydrift
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DIRECTIONS FOR USE
Table 1: Hand Application
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

Situation	Weeds Controlled	State	Rate g/m²	Critical Comments
Brigalow regrowth and associated problem woody weeds on grazing land	Belah	Qld only	1.5	Estimate the area within 30 cm beyond the drip zone of each target weed or group of weeds and calculate the amount of Albaugh TEBIX 200 GR Herbicide to cover area to be treated. Distribute the required dose uniformly within this area. Use the higher rate for heavy density regrowth and heavy clay soils. [1] Some suppression of these species is provided by the rates given. [2] DO NOT use within 30 m of trees. DO NOT apply to single continuous area greater than 0.5 ha in size.
	Brigalow, Currant Bush	NSW & Qld only	1 to 1.5	
	Dawson Gum	Qld only	1	
	False Sandalwood [1]	NSW & Qld only	2	
	Holly Bush		1.5	
	Limebush		1	
	Poplar Box	Qld only		
	Whitewood	NSW, Qld & NT only		
	Yellowwood	Qld only		
Tea tree regrowth and associated woody weeds on grazing land	Black Tea Tree	Qld, WA & NT only	1.5	
	Broadleaf Tea Tree		1	
	Cocky Apple [1]		2	
	Paperbark Tea Tree, Poplar Gum		1.5	
	Swamp Box		2	
Other woody weeds on grazing land	African Boxthorn	NSW, Qld & NT only	2	
	Blue Heliotrope [2]	NSW only	0.5	
	Broadleaf Ironbark, Brown Box	Qld only	1.5	
	Coolibah	Qld & NT only	1	
	Gidgee	Qld only	0.5 to 1	
	Gum-topped Box		1.5	
	Groundsel Bush	NSW & Qld only	1	
	Parkinsonia	Qld & NT only	1 to 1.5	
	Prickly Acacia, Rubbervine	Qld & NT only	1.5	
	Wild Rosemary	NSW & Qld only		
Giant sensitive plant on grazing land and other grasslands	*Mimosa pigra*	NT only	1	Apply after the onset of storms (to moisten soil, close soil cracks (where applicable) and freshen growth of giant sensitive tree) and before wet season surface flooding. Repeat application may be required in year two for long term control. Some grass suppression may occur.

DIRECTIONS FOR USE: Aerial and Ground Application
See GENERAL INSTRUCTIONS - APPLICATION METHODS Section for application method details.

Situation	Weeds Controlled	State	Rate kg/ha	Critical Comments
Brigalow regrowth and associated problem woody weeds on grazing land	Belah, Brigalow, Currant Bush, Dawson Gum, Limebush, Poplar Box, Scrub Boonaree, Whitewood, Wild Rosemary, Yellowwood	Qld only	7.5 to 15	For brigalow areas, a guide to the recommended rates is 7.5 kg for lighter duplex soils, 10 kg for medium density regrowth on light clay soils, 12.5 kg and above for heavy density regrowth on heavy acidic clays.
	False Sandalwood		10 to 15	On False Sandalwood partial control and suppression only will be obtained at the specified rate.
Tea tree regrowth and associated woody weeds on grazing land	Broadleaf Tea Tree, Paperbark Tea Tree		5 to 7.5	For tea tree use the highest rate if the area is prone to waterlogging. Excessive rainfall or waterlogging may reduce control.
	Cocky Apple*, Swamp Box		10	* For Cocky Apple partial control and suppression only will be given at the specified rate.
Other woody weeds on grazing land	Gidgee		5 to 10	Use the high rate on dense growth or heavy clay soils.
	Parkinsonia	Qld & NT only	10 to 15	
	Prickly Acacia			
	Poplar Gum		7.5 to 10	
	Rubbervine		7.5 to 15	
Eucalypt woodland management using Property Management Planning Principles	Poplar Box	Qld only	7.5 to 10	Use the higher rate on dense growth or heavier clay soils. To be used in accordance with any "Tree Clearing Guidelines" and 20% intact retention of species.
Giant sensitive plant on grazing land and other grasslands	*Mimosa pigra*	NT only	5 to 10	Apply after the onset of storms (to moisten soil, close soil cracks (where applicable) and freshen growth of giant sensitive tree) and before wet season surface flooding. Repeat application may be required in year two for long term control. Some grass suppression may occur.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

FOR NATIVE VEGETATION
USE OF ALBAUGH TEBIX 200 GR HERBICIDE ON NATIVE VEGETATION MUST BE DONE IN ACCORDANCE WITH STATE AND/OR LOCAL LEGISLATION.
WITHHOLDING PERIODS
NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS
Review the restraints before reading these instructions.
Albaugh TEBIX 200 GR Herbicide may be applied at any time of the year. Treatments become effective after sufficient rainfall has occurred to move the chemical into the root zone where it is absorbed by woody plants. Applications just prior to seasonal rainfall give the most rapid response.
- Herbicidal activity will continue for several seasons after application. During this time woody plants may go through repeated defoliations and regrowths until death occurs. A single application is normally effective for several years.
- Forage grass production usually increases as woody weed competition is reduced. Increased grass production is also dependent on adequate rainfall and an acceptable grazing management program.
- It is important to allow grasses to seed and seedlings to become established after treatment before grazing pressure is increased. Areas treated with Albaugh TEBIX 200 GR Herbicide may be overseeded with suitable pasture species.
- Consult your agricultural specialist for details on suitable seeding rates, timing and fertiliser application program.
- The residual action of Albaugh TEBIX 200 GR Herbicide may be impaired if fire occurs before clay pellets are dissolved and the chemical has moved into the root zone.

APPLICATION METHODS
HAND APPLICATION
Gloves must be used during hand application. Apply Albaugh TEBIX 200 GR Herbicide by uniformly distributing by hand over the desired area after weighing out the recommended quantity of pellets. The best way to calibrate yourself is to weigh a normal handful of pellets. If, for example, a handful weighs 20 grams, then this will treat 10 to 20 m^2 at the rate of 0.5 to 1 g/m^2.
DISPENSER PACK
Apply Albaugh TEBIX 200 GR Herbicide by uniformly distributing by hand after dispensing a single dose (7 g) from the dispenser unit. Apply the pellets at 1 g to 2 g per metre square. One dose (7 g) will treat 7 to 3.5 square metres at these rates.
ADDITIONAL INSTRUCTIONS
When applying pellets, distribute them under the tree or regrowth canopy from the root crown or stems to 20 cm beyond the drip zone of the plants you want to control. The time required for complete plant death depends on soil type, amount of rainfall, root depth, plant species, plant density and rate of Albaugh TEBIX 200 GR Herbicide used. Some species may go through several defoliations from 6 months to 2 years before dying. Timber and regrowth control with Albaugh TEBIX 200 GR Herbicide pellets may be variable when used on soils with high clay content, high organic matter or when treating deep rooted or more tolerant species.
TREATING INDIVIDUAL TREES BY HAND
Estimate the area within 30 cm beyond the drip zone, i.e. the area under the plant canopy. It is essential that this area is treated as evenly as possible with pellets being spread from the tree trunk to slightly beyond the drip line formed by the branches.
Do not throw all the pellets at the base of the tree trunk. This will waste material and could result in poor control.
TREATMENT OF CLUMPS OF TREES OR PATCHES OF REGROWTH BY HAND
Albaugh TEBIX 200 GR Herbicide pellets are particularly effective in treating areas of regrowth. Estimate the area within 30 cm beyond the drip zone and apply the Albaugh TEBIX 200 GR Herbicide pellets on a grid pattern to uniformly cover the area. Pellets should be applied over the top of regrowth if at all possible in an attempt to maintain a good distribution pattern.
TREATING FENCE LINES BY HAND
Estimate the area to be treated, along the fence line and then apply Albaugh TEBIX 200 GR Herbicide pellets on both sides of the fence line as evenly as possible within the area extending 30 cm out beyond drip zone.
AERIAL AND GROUND APPLICATION
Apply Albaugh TEBIX 200 GR Herbicide by uniformly distributing with an approved aircraft or ground driven air equipment, after calibration to achieve the desired application rate. Albaugh TEBIX 200 GR Herbicide may be applied at any time of year. Treatments become effective after sufficient rainfall has occurred to move the chemical into the root zone where it is absorbed by woody plants.
Applications just prior to seasonal rainfall give the most rapid response. Herbicidal activity will continue for several seasons after application. During this time woody plants may go through repeated defoliations and re-growth until death occurs. A single application is normally effective for several years. Forage grass production usually increases as woody weed competition is reduced. Increased grass production is also dependent on adequate rainfall and an acceptable grazing management program. It is important to allow grasses to seed and seedlings to become established after treatment before grazing pressure is increased. Areas treated with Albaugh TEBIX 200 GR Herbicide may be over seeded with suitable pasture species. Consult your agricultural specialist for details of suitable seeding rates, timing and fertilizer application program. The residual action of Albaugh TEBIX 200 GR Herbicide may be impaired if fire occurs before clay pellets are dissolved and the chemical has moved into the root zone.
1. Aerial Application
Aerial application of Albaugh TEBIX 200 GR Herbicide can be carried out by accredited fixed wing aircraft or helicopters fitted with specially designed application apparatus. Use a Positive Metering Device (Meterate) for fixed wing aircraft and Isolair or similar precision equipment for helicopters.
A minimum flying height of 20 metres above the canopy of the target plants must be maintained. The swath width is dependent on the type of spreader used. The target areas must be correctly marked out and flown with the aid of GPS (Global Positioning System) or markers clearly visible to the operator or a method of marking approved by ALBAUGH AUSTRALIA PTY LTD.
2. Ground Application
Ground applications of Albaugh TEBIX 200 GR Herbicide can be carried out in suitable areas by accredited operators.

CLEANING APPLICATION EQUIPMENT
Thoroughly clean all traces of Albaugh TEBIX 200 GR Herbicide from vessels holding pellets e.g. bucket and gloves after use. Residues cleaned from this application equipment should not be emptied on areas where they will contact roots of desirable trees, shrubs or plants.
RESISTANT WEEDS WARNING
GROUP 5 HERBICIDE
Albaugh TEBIX 200 GR Herbicide is a member of the Ureas group of herbicides. The product has inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management, the product is a Group 5 herbicide.
Some naturally occurring weed biotypes resistant to the product and other Group 5 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 5 herbicides.
Since the occurrence of resistant weeds is difficult to detect prior to use, ALBAUGH AUSTRALIA PTY LTD accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture or local ALBAUGH AUSTRALIA PTY LTD representative.
PROTECTIONS OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS
DO NOT allow any roots of desirable plants, trees or ornamentals to contact Albaugh TEBIX 200 GR Herbicide. Albaugh TEBIX 200 GR Herbicide may cause temporary herbicidal symptoms to appear on perennial grasses. Higher rates of application of Albaugh TEBIX 200 GR Herbicide (2 g/m^2) may kill grasses, particularly native annuals. This effect can be minimised by application when grasses are dormant.
Albaugh TEBIX 200 GR Herbicide may seriously injure desirable forage plants such as pasture legumes and other broadleaf plants. These species may regenerate in subsequent seasons.
PROTECTION OF LIVESTOCK
Poisonous plants may become more palatable after application of granules and stock should be kept away from these plants until they have died down.
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
DO NOT contaminate streams, rivers or watercourses with the granules or used containers.
STORAGE AND DISPOSAL
Store in the stored original container in a dry, cool, well-ventilated area out of direct sunlight. DO NOT store near food, feedstuffs, fertilisers or seed. DO NOT dispose of granules on-site.
Single-rinse or shake remainder into the application equipment until the bag is empty. DO NOT dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.
SMALL SPILL MANAGEMENT
Sweep up material for disposal and contain in a refuse vessel for disposal (see Storage and Disposal Section).
SAFETY DIRECTIONS
Harmful if swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing the product, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.
FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.
SAFETY DATA SHEET
Additional information is listed in the safety data sheet (SDS) which is available from the supplier and accessible from the Albaugh website albaugh.com/au
TERMS AND CONDITIONS OF SUPPLY, SALE AND USE
Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of strains resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but we cannot exercise any control over their mixing, use or application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application, or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

APVMA Approval Number: 94743/163025

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture:
Harmful if swallowed
Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. IF SWALLOWED: Call a POISON CENTRE/doctor if you feel unwell. Rinse mouth. Dispose of contents/ container in accordance with local regulations.

Batch No:

Date of Manufacture:

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

ALBAUGH TEBIX 200 GR HERBICIDE

Face

Live Copy Area

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ALBAUGH TEBIX 200 GR HERBICIDE

ACTIVE CONSTITUENT: 200 g/kg TEBUTHIURON

GROUP 5 HERBICIDE

For control of brigalow regrowth, tea tree regrowth, Mimosa pigra and certain problem woody weeds on grazing lands by hand, aerial and ground application as specified in the Directions for Use table.
IMPORTANT: READ THE LABEL INSTRUCTIONS AND DIRECTIONS FOR USE BEFORE OPENING AND USING.

Net Contents
15 KG

SELECTIVE HERBICIDE

ALBAUGH AUSTRALIA PTY LTD
Level 1, 530 Little Collins Street,
Melbourne VIC 3000
Australia
Tel 03 9909 7183
www.albaugh.com/au

ALBAUGH® your alternative

ALBAUGH TEBIX 200 GR HERBICIDE

PBOM 11x5.5x29 .067

PMS 371 PMS 5493

Tebix 200 GR_15kg