

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with ALBAUGH COVERPRO 480 SC Fungicide. If you are growing produce for export, please check with supplier for the latest information on MRLs and import tolerances before using ALBAUGH COVERPRO 480 SC Fungicide.

GENERAL INSTRUCTIONS

Foliar diseases on cereal crops

Monitor the crop regularly for symptoms of disease. Generally spray at the first sign of disease, although this will depend on factors such as expected weather conditions and the particular crop variety resistance. Refer to Directions for Use for particular disease recommendations. Up to two sprays of Albaugh COVERPRO 480 SC may be applied per season to the crop. Ensure good coverage of all susceptible plant parts.

Disease control in oats

Caution:

Varieties most at risk may exhibit early senescing and bronzing of leaves trait under various stress conditions not related to fungicide sprays.

Disease control in canola

Blackleg

Higher blackleg risk can be expected in higher rainfall districts (above 500 mm annual rainfall), where crops are grown within 500 m of a previous year’s stubble and in later sown crops (May to August). Other factors will also increase the risk of blackleg infection, including the intensity of canola cropping in a district, rainfall before sowing and the frequency of growing the same canola cultivar. Consult industry guidelines for more detailed assessment of blackleg risk in specific situations. Up to two sprays of Albaugh COVERPRO 480 SC may be applied per season to the crop.

Sclerotinia

Albaugh COVERPRO 480 SC is most effective when application is made prior to conditions conducive to sclerotinia infection. Infection and disease development are most conducive in warmer winter or spring conditions with extended periods of leaf wetness due to rainfall, dew and high humidity. Sclerotinia is most likely to develop where day temperatures are warmer coinciding with a saturated soil profile and rainfall events. Refer also to industry guidelines for advice on conditions under which sclerotinia are most likely to develop. Control of sclerotinia stem rot is more effective in crops which have a uniform flowering. Uneven flowering (e.g. caused by staggered germinations) makes optimum spray timing difficult and two sprays may be required in these crops.

Generally a single application of Albaugh COVERPRO 480 SC at 20 to 30% flowering will control sclerotinia in crops with a short flowering interval. Crops with an extended flowering period may require a second application prior to 50% flowering (full-bloom) to adequately control sclerotinia if conditions late in the season are conducive to development of disease. Length of protection may be reduced in bulky crops where coverage is difficult and where there is growth dilution of the fungicide. For optimum protection, applications should be directed to obtain coverage on petals, leaves and stems. **MIXING** Prior to pouring, shake container vigorously, then add the required quantity of Albaugh COVERPRO 480 SC to water in the spray vat with agitators in motion. Add the required amount of adjuvant if necessary and mix thoroughly.

APPLICATION

Ground:

Wheat, barley, and oats: Apply product using a spray volume of 70 – 100 L/ha and a MEDIUM spray droplet size category.

Canola: Apply product using a spray volume of 60 – 100 L/ha and a MEDIUM pray droplet size category.

COMPATIBILITY

USE OF ADJUVANT

Suitable Adjuvants	Comments
non-ionic surfactant at 0.25% v/v	Can be used at all rates of Albaugh COVERPRO 480 SC for ground application.

RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Albaugh COVERPRO 480 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Albaugh Australia Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

Re-entry Period

Do not enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves. Clothing must be laundered after each day’s use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. Do not use empty container for any other purpose.

SAFETY DIRECTIONS

May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. Wash hands after use. After each day’s use, wash gloves and contaminated clothing.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS) which is available from the supplier and accessible from the Albaugh website albaugh.com/au

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of strains resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but we cannot exercise any control over their mixing, use or application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application, or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

APVMA Approval Number: 95990/147834