

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ACTIVE CONSTITUENT: 480 g/L METRIBUZIN

For selective weed control in various broadacre and vegetable crops
IMPORTANT: READ THE ATTACHED LEAFLET/BOOKLET BEFORE USE

ALBAUGH AUSTRALIA PTY LTD
Level 1, 530 Little Collins Street,
Melbourne VIC 3000
Australia
Tel 03 9909 7183
www.albaugh.com/au



DIRECTIONS FOR USE

CROP/ SITUATION	WEEDS	STATE	RATE	CRITICAL COMMENTS
Asparagus	False castor oil (thornapples), shepherd's purse, fat hen Winter weed control: Chickweed, shepherd's purse, nettle, deadnettle Spring and summer weed control: False castor oil (thornapples), shepherd's purse, fat hen	Qld, NSW, ACT, Vic, Tas, SA only WA only	1.1 or 2.2 L/ha 0.8 or 1.5 L/ha	Direct seeded asparagus Apply to moist soil as soon as possible after sowing before crop emerges and overhead irrigate. A follow up application may be required 4-8 weeks later for emerged weeds. DO NOT spray until asparagus seedlings are at least 4 cm high. Some temporary burning may occur. Use the higher rate on heavier soils. Seedling asparagus (transplanted) DO NOT spray before asparagus seedlings are 4 cm high and seedlings have been well established after transplanting. Spray when weeds have emerged when young and actively growing. Use the higher rate on older weeds. Established asparagus in beds Apply immediately after transplanting crowns or preparing beds for cutting and before spears emerge, or after cutting for the season is finished. Overhead irrigate within a few days after spraying. Use the higher rate on heavier soils.
Barley Post-emergent application	Controlled: seedlings of: capeweed, cats ear, charlock, chickweed, common cotula, corn gromwell (sheepweed, white ironweed), corn spurry, creeping speedwell, crown beard (dogweed), deadnettle, dock, fat hen, field madder, fumitory, hare's ear (treadle mustard), heliotrope, hogweed (wireweed), horehound, Indian hedge mustard, lesser swinecress, mouse-ear chickweed, mountain sorrel (purple calandrinia), rough poppy, scarlet pimpernel, scrub nettle (stinging nettle), shepherd's purse, sorrel, sowthistle, spiny emex (threecornered jacks), storksbill, toad rush, wild radish, wild turnip, winter grass, yellow burnweed Suppressed: Annual ryegrass, sub clover Buxbaum's speedwell, Powell's amaranth, stagger weed, sun spurge, great brome Buchan weed, fog grass Toad rush	Vic, Tas, SA only Tas only SA only NSW, ACT, Vic, SA only	280 mL/ha (Light sandy soils i.e. sandy loam to loamy sands) 435 mL/ha (Medium soils, loams. Silt plus clay content 40 to 60%) 580 mL/ha (Heavy soils, clay loams. Silt plus clay content over 60%) 150 mL	DO NOT spray before barley is at 3 leaf stage or later than 8 weeks after sowing. Application of this product on alkaline sandy soils (pH greater than 7) will result in severe crop damage. In Tasmania , DO NOT use Albaugh Rometri 480 SC Selective Herbicide on barley grown on sandy soils. For effective wild radish, wireweed and ryegrass control, the weed must have emerged before Albaugh Rometri 480 SC Selective Herbicide application. Wireweed in the cotyledon stage and annual ryegrass with no more than 2 leaves can be controlled provided rain follows within a few days of spraying. Emerged weeds must be young and actively growing. Toad rush should be sprayed at the 2-4 leaf stage for optimum control. Spray after rain when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on soil surface.
Barley (except Morrell and Harrington) Pre-sowing or incorporated by sowing	Capeweed, doublegee, geranium, mustard, toad rush, turnip, wild radish, wireweed Suppressed: Annual ryegrass, barley grass, brome grass	WA - Great Northern region only	210 - 235 mL/ha + 1 L/ha Trifluralin	1. Correct application and incorporation techniques are critical to obtain optimum performance of Albaugh Rometri 480 SC Selective Herbicide and Trifluralin mixtures. ENSURE all recommendations on the Trifluralin label are followed. 2. Use the lower rate in lighter soils, and the higher rate in heavier soils. 3. Application of this product on alkaline sandy soils (pH greater than 7) will result in severe crop damage. 4. Apply to moist soil for best efficacy. 5. Seed below herbicide band at 5 cm. 6. Reduced control may occur under high weed burdens (> 100/m ²). 7. The performance of Albaugh Rometri 480 SC Selective Herbicide and Trifluralin mixtures in terms of crop tolerance and weed control may vary according to climate, soil and crop conditions. This treatment should only be used where no alternatives are available.

CROP/ SITUATION	WEEDS	STATE	RATE	CRITICAL COMMENTS
Chickpeas	Controlled: as per Barley (Vic, SA, Tas) Buchan weed, fog grass (SA only) Suppressed: Sub clover	Southern NSW, ACT, Vic, <u>SA, WA only</u>	280 mL/ha (Light sandy soils i.e. sandy loam to loamy sands)	ONLY apply post sowing, pre-emergence. Crops should be sown at least 5 cm deep. DO NOT apply post-emergence, as crop injury will occur. DO NOT apply to chickpeas grown on sandy soils, i.e. silt plus clay less than 40%. The vigour of lentils may be reduced, especially if heavy rain falls after spraying. Weeds should be from pre-emergence to 3 leaf stage except wireweed (hogweed) which should not be beyond the cotyledon stage. Best results for grass and wireweed control are obtained when rain or irrigation follows within a few days of spraying. DO NOT apply by aircraft to chickpeas and faba beans.
Faba beans		NSW, ACT, Vic, <u>SA, WA only</u>		
Lentils		Vic, SA, WA only	435 mL/ha (Medium soils, loams. Silt plus clay content 40 to 60%) 580 mL/ha (Heavy soils, clay loams. Silt plus clay content over 60%)	
Lupins (except Danja, Kiev Mutant, Tanjil, Wonga, Belara, Chitick, Moonah, Quilinoack, Woodjil or Albus lupins)	For control of wild radish up to 250 mm diameter. For suppression of capeweed, doublegee, and larger wild radish. Use higher rate for larger weeds.	WA – Great Northern region ONLY	160 to 235 mL/ ha plus 100 mL Brodal® Options	1. Albaugh Rometri 480 SC Selective Herbicide should only be used following a pre sowing treatment of Simazine. 2. DO NOT apply to alkaline soils with pH > 7. 3. Apply to actively growing lupins from the 3-4 leaf stage onwards. 4. DO NOT apply to weeds or crop which are under stress. 5. Apply to moist soil for best efficacy. 6. Use higher rates for heavy weed burden and larger weed size. Apply before weeds are covered by crop canopy. 7. Best results are obtained if good soil moisture exists at, and received soon after application. 8. DO NOT apply wetting agents, spray oils or grass selective herbicides with the Albaugh Rometri 480 SC Selective Herbicide + Brodal Options mixture. 9. Transient crop yellowing and burning may become evident after application. This effect is short term and will not normally reduce lupin yield. 10. Refer to Critical Comments section on the Brodal Options label. IMPORTANT The performance of Albaugh Rometri 480 SC Selective Herbicide in terms of crop tolerance and weed control may vary with varying climate, soil and crop conditions. This treatment should only be used where no alternatives are available.
Oats	Toad rush	Vic only	150 mL/ha	Toad rush should be sprayed at the 2-4 leaf stage for optimum control. Spray after rain when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on soil surface.
Peas (field, processing, green) DO NOT use on Laura, Pennant or White Brunswick varieties	Controlled: as per Barley (Vic, SA, Tas) Suppressed: Annual ryegrass, sub clover Buxbaum's speedwell, Powell's amaranth, stagger weed, sun spurge Buchan weed, fog grass	Qld, NSW, ACT, Vic, Tas, SA only Tas only SA only	280 mL/ha (Light sandy soils i.e. sandy loam to loamy sands) 435 mL/ha (Medium soils, loams. Silt plus clay content 40 to 60%) 580 mL/ha (Heavy soils, clay loams. Silt plus clay content over 60%) 730 mL/ha (Koo-wee-rup swamp and volcanic soils)	Pea crops should be sprayed pre-emergence or early post-emergence of crop and/or weeds, but within 6 weeks after sowing (and not later than 8 weeks in Victorian winter crops). Peas must not be beyond the 3 node stage. DO NOT treat peas post-emergence in areas or soils where collar rot, stem rot or black spot are present or are likely to occur. In Tasmania, DO NOT use Albaugh Rometri 480 SC Selective Herbicide on field peas grown on light sandy soils. DO NOT spray Cressy Blue peas post-emergence. Some other varieties, e.g. Dundale and Wirrega, may be especially sensitive to Albaugh Rometri 480 SC Selective Herbicide. DO NOT use on a large scale unless it has been ascertained that the variety shows sufficient tolerance under local conditions. For effective wild radish, wireweed and ryegrass control, the weed must have emerged before Albaugh Rometri 480 SC Selective Herbicide application. Wireweed in the cotyledon stage and annual ryegrass with no more than 2 leaves can be controlled provided rain follows within a few days of spraying. Emerged weeds must be young and actively growing.
Pigeon peas	Apple of Peru, bladder ketmia, gooseberry, Noogoora burr, cowvine (peach vine), volunteer cotton, yellow vine, false castor oil (thornapples)	Qld, NSW, ACT only - except sandy soils	750 mL/ha	Rain grown and overhead irrigated crops: Apply immediately after or within a day or two after sowing before crops emerge then irrigate. Otherwise rain after spraying before weeds emerge is necessary to get useful weed control. The longer the time interval between spraying and rain or irrigation the less the effectiveness of the herbicide. Albaugh Rometri 480 SC Selective Herbicide can be applied to a band centred on crop rows for in-row weed control. Untreated soil between treated bands thrown onto treated areas during subsequent cultivation will reduce weed control along rows. Albaugh Rometri 480 SC Selective Herbicide can be applied after pre-sowing Trifluralin treatment. Avoid leaving a depression over the row during planting to enhance crop tolerance. Furrow irrigated crops: Apply after furrowing out and within two weeks before sowing and incorporate with Lillistone cultivators to finely tilled soil free of emerged weeds. Where grass weeds are expected, add Trifluralin at the rate recommended by the manufacturer for the soil type. For post-emergence weed control: Apply when weeds are in the young seedling stage and actively growing, provided crop plants have at least 2-trifoliolate leaves. DO NOT spray if rain is likely to fall within several hours. Transitory damage may occur to the pigeon pea crop but yields will not be significantly affected.
Potatoes Some varieties such as Atlantic and red skin varieties are especially sensitive to Albaugh Rometri 480 SC Selective Herbicide. Damage may occur even at the lowest label rate. DO NOT use on Kurell or Bison potatoes.	Controlled: as per Barley (Vic, SA, Tas) Suppressed: Annual ryegrass, dock, sub clover, summer grasses (<i>Digitaria</i> spp.)	All States	730 mL/ha or 1.1 L/ha	Spray when potatoes are emerging with the largest shoots not more than 10 cm tall. Use higher rate for emerging summer grasses. In summer crops on soils which dry rapidly on the surface, the young actively growing weeds will be killed but residual effect may be reduced. If there is a subsequent germination of weeds apply a second spray at 730 mL/ha (except NSW). DO NOT use Albaugh Rometri 480 SC Selective Herbicide in Western Australia in the Perth metropolitan area; on the south west winter potato crops; or on Delaware potatoes on light soils.
		Vic - Bellarine Peninsula only	430 to 580 mL/ha (winter)	Potatoes may be hilled before or after spraying Albaugh Rometri 480 SC Selective Herbicide. If hilled after spraying a good rain or irrigation is desirable between spraying and hilling. No other cultivation should be necessary. Use higher rate for emerging summer grasses.
		Qld - Atherton Tableland ONLY	730 mL/ha for winter crops 1.45 L/ha for summer crops	Hill up the crop soon after planting. Irrigate to allow weed seeds to germinate and potatoes to begin emerging. Apply as an overall spray to a soil moist to the surface, before all the crop has emerged.

CROP/ SITUATION	WEEDS	STATE	RATE	CRITICAL COMMENTS
Soybeans in soils of pH 9 or lower DO NOT use on Cannapolis, Hill, Semstar or Triton varieties	Apple of Peru, bladder ketmia, cowvine (peach vine), gooseberry, Noogoora burr, volunteer cotton	NSW - clay soils in NW only, Qld - clay soils on Darling Downs only	750 mL/ha	Rain grown and overhead irrigated crops: Apply immediately after or within a day or two after sowing before crops emerge then irrigate. Otherwise rain after spraying before weeds emerge is necessary to get useful weed control. The longer the time interval between spraying and rain or irrigation the less the effectiveness of the herbicide. Albaugh Rometri 480 SC Selective Herbicide can be applied to a band centred on crop rows for in-row weed control. Untreated soil between treated bands thrown onto treated areas during subsequent cultivation will reduce weed control along rows. Albaugh Rometri 480 SC Selective Herbicide can be applied after pre-sowing Trifluralin treatment. Avoid leaving a depression over the row during planting to enhance crop tolerance. Furrow irrigated crops: Apply after furrowing out and within two weeks before sowing and incorporate with Lillistone cultivators to finely tilled soil free of emerged weeds. Where grass weeds are expected, add Trifluralin at the rate recommended by the manufacturer for the soil type. DO NOT use on cv Triton as crop damage may occur.
Tomatoes	Controlled: As per Barley (Vic, SA, Tas) plus barnyard grass, bull's head, crowsfoot grass, nightshade, pigweed, black pigweed, summer grass, green summer grass	Qld, NT only	1.45 L/ha (15 mL/100 m of row)	Apply in a 1 metre wide band centred on the inter-row either during or shortly after the formation of the final irrigation furrow. Apply to soil moist to the surface. Apply through applicators fitted with flat fan nozzles and operated at pressures not exceeding 200 kPa to avoid misting and spray drift. DO NOT follow autumn plantings of tomatoes with cucurbits or capsicums in the same season.
Transplanted tomatoes	Apple of Peru, potato weed	NSW, ACT only	750 mL - 1.2 L/ha	Apply as a directed spray to weeds at the base of tomato plants. Use the lower rate when tomatoes have 4-5 expanded leaves or the higher rate when tomatoes beyond this stage and weeds are well established. Accurately calibrated spray equipment, fitted with one flat fan 80° even spray Teejet nozzle operated to deliver 100 to 200 L of spray per hectare and angled to avoid tomato leaves should be used. DO NOT apply during or soon after prolonged periods of overcast or cloudy weather. Allow at least two fine, sunny days before spraying. Spray on a sunny fine day. Suppression of other broadleaf weeds such as false castor oil may occur.
Vetch	Controlled: as per Barley (Vic, SA, Tas)	Vic, SA, WA only	280 mL/ha (Light sandy soils i.e. sandy loam to loamy sands) 435 mL/ha (Medium soils, loams. Silt plus clay content 40 to 60%) 580 mL/ha (Heavy soils, clay loams. Silt plus clay content over 60%)	Crop: Only apply post sowing pre-emergence. Crop should be sown at least 5 cm deep. The vigour of vetch may be reduced, especially if heavy rain falls after spraying. Do not apply post-emergence, as crop injury will result. Weeds: Weeds should be from pre-emergence to 3 leaf stage except wireweed (hogweed) which should not be beyond the cotyledon stage. Best results for grass and wireweed control are obtained when rain or irrigation follows within a few days of spraying.
Wheat Post-emergent application only	Toad rush	NSW, ACT, Vic, SA only	150 mL/ha	Toad rush should be sprayed at the 2-4 leaf stage for optimum control. Spray after rain when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on soil surface.
Wheat (Blade & Eagle Rock only) Pre-sowing or incorporated by sowing	Capeweed, common cotula, doublegee, fumitory, Indian hedge mustard, toad rush, wild turnip, wild radish, wireweed Suppressed: Brome grass (<i>Bromus diandrus</i> & <i>B. rigidus</i>)	WA – Great Northern region only Sandy – sandy loam soils	310 mL/ha	1. Albaugh Rometri 480 SC Selective Herbicide has increased crop safety and gives more reliable weed control when incorporated by sowing and trailing harrows. 2. DO NOT apply to alkaline soils with pH > 7. 3. Apply to moist soil for best efficacy. 4. Seed below herbicide band at 5 cm.
	Annual ryegrass, barley grass, geranium		310 mL/ha + 1 L/ha Trifluralin	5. Reduced control may occur under high weed burdens (> 100/m²). 6. To improve control of annual ryegrass and barley grass add Trifluralin at 1 L/ha. 7. Ensure Albaugh Rometri 480 SC Selective Herbicide is fully dispersed in the boom spray tank prior to Trifluralin addition. 8. The performance of Albaugh Rometri 480 SC Selective Herbicide and Trifluralin mixtures in terms of crop tolerance and weed control may vary according to climate, soil and crop conditions. This treatment should only be used where no alternatives are available. 9. Refer to the Application section of this and the Trifluralin label for complete instructions on product use.
White Lupins	Buchan weed, capeweed, charlock, Chickweed, Deadnettle (<i>Lamium</i> spp.) fat hen, fumitory (common and wall fumitory – Tas), wireweed (hogweed), lesser swinecress, purple calandrinia, scarlet pimpernel, spiny emex, corn spurry (spurry - Tas), stinging nettle (dwarf nettle - Tas), toad rush, winter grass, heliotrope, wild oats, wild radish, wild turnip Suppression only: Sub clover and docks (broadleaf dock - Tas)	Tas only	625 mL/ha	Spring application. Apply after sowing but before crop emerges.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

Withholding Periods:

Harvest

Asparagus: **DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

Tomatoes: **DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION**

Other crops: **NOT REQUIRED WHEN USED AS DIRECTED**

Grazing

All uses: **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

Trade Advice: Export of Treated Produce

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Albaugh Rometri 480 SC Selective Herbicide. If you are growing edible produce for export, please check with Albaugh Australia Pty Ltd for the latest information on MRLs and import tolerances before using Albaugh Rometri 480 SC Selective Herbicide.

General Instructions

Albaugh Rometri 480 SC Selective Herbicide is effective against a range of broadleaf weeds and some grasses. Albaugh Rometri 480 SC Selective Herbicide is absorbed by roots, shoots and leaves of weeds. It can therefore be applied before or after weeds emerge depending on the tolerance of the crop being treated. For recently introduced varieties DO NOT use Albaugh Rometri 480 SC Selective Herbicide unless it is ascertained that the variety to be treated shows sufficient tolerance to the treatment under local climatic conditions.

CROP SAFETY

The use of other herbicides and/or crop oils on broadleaf crops treated with Albaugh Rometri 480 SC Selective Herbicide may cause severe damage. Before using Albaugh Rometri 480 SC Selective Herbicide on newly released crop varieties, confirm the variety tolerance with the State Department of Agriculture, Agronomist or Albaugh Australia Pty Ltd representative.

MIXING

Prior to pouring, shake container vigorously for at least 30 seconds, then add the required quantity of Albaugh Rometri 480 SC Selective Herbicide to the required amount of water and stir. Rinse empty container with water and pour into spray tank. Keep agitators in motion.

APPLICATION

General

Best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or sprinkler irrigation (6-12 mm) within 2 weeks after application. If heavy rains occur soon after application, injury to the crop may result.

Pre-emergence

Soil should preferably be moist to the surface and must not be cloddy or have residues of a previous crop on the surface. It is better to delay pre-emergence spraying of loose or fluffy soils until there is a good soil settling rain. Rain or irrigation soon after spraying is required to activate Albaugh Rometri 480 SC Selective Herbicide.

Post-emergence

Post-emergence spraying will control young actively growing emerged weeds through leaf uptake. Subsequent pre-emergence control of later germinating weeds requires rain or irrigation after spraying. Crops may show slight chlorosis or necrosis after spraying. Two sunny days before spraying enhances crop tolerance. Heavy rain (80-100 mm) after spraying on deep sandy or friable well structured soils may leach Albaugh Rometri 480 SC Selective Herbicide beyond the root zone thus reducing residual effect.

Ground Spraying

Apply Albaugh Rometri 480 SC Selective Herbicide through a carefully calibrated boom spray set at 200 to 300 kPa pressure (except for tomatoes in Qld only) fitted with fan nozzles. Use a convenient quantity of water (50-100 L/ha). Cone nozzles and other means of application are not satisfactory and should not be used. When spraying avoid over-lapping and shut off spray booms while starting, turning slowly and when stopped. Neglecting these procedures will cause higher rates of application resulting in crop damage. Delay post-emergence spraying if rain is expected within a few hours.

Aerial Spraying

Wind: Spray only in crosswinds, otherwise uneven coverage may result. DO NOT spray onto inversions or in dead calm conditions. DO NOT spray when winds are light and variable in direction unless smoke indicators or the spray cloud show that the spray is going down into the crop. Spray drift may damage adjoining crops. Make sure that the wind is blowing away from crops for which Albaugh Rometri 480 SC Selective Herbicide is not recommended.

Flying: Fly with the wheels 2 to 3 metres above the crop as “wheels on crop” spraying causes uneven distribution. Track spacing must be uniform and suited to the type of aircraft and crosswind speed.

Droplet Size: For grasses and vertical weeds, use fine droplets. For horizontal weeds and drift limitation, use droplets over 200 microns and increase volume.

Water Volume: Use 20 to 30 L water/ha.

RESISTANT WEEDS WARNING

GROUP	5	HERBICIDE
-------	---	-----------

Albaugh Rometri 480 SC Selective Herbicide is a member of the triazinone group of herbicides and has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management the product is a Group 5 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 5 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 5 herbicides. Annual ryegrass biotypes resistant to diclofop-methyl and other 'grass specific' herbicides are often also resistant to Albaugh Rometri 480 SC Selective Herbicide. Before using Albaugh Rometri 480 SC Selective Herbicide on a population resistant to ‘grass specific’ herbicides, have a resistance test conducted to ensure that it is still susceptible to Albaugh Rometri 480 SC Selective Herbicide.

Since the occurrence of resistant weeds is difficult to detect prior to use, Albaugh Australia Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the survivors. DO NOT make more than one application of a Group 5 herbicide to a crop in the same season, either pre-sowing incorporated by sowing or post crop and weed emergence. If the user suspects that a Group 5 resistant weed is present, Albaugh Rometri 480 SC Selective Herbicide or other Group 5 herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Albaugh Australia Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND NON-TARGET PLANTS

Prevent drift of spray to sensitive plants. Keep from contact with fertilisers, insecticides, fungicides and seeds. Thoroughly clean all traces of Albaugh Rometri 480 SC Selective Herbicide from application equipment immediately after use. To minimise risk to follow-on sensitive crops such as beetroot, brassicas, capsicums, cotton, cucurbits, lettuce, onions and sunflowers, it is advisable to work land thoroughly - allow at least 6 months (12 months for cucurbits) between application of Albaugh Rometri 480 SC Selective Herbicide and sowing seed of these sensitive crops. A longer period may be required following long dry periods between crops. If rates of Albaugh Rometri 480 SC Selective Herbicide higher than 1.5 L/ha were used, allow at least 12 months before planting these crops. The use of other herbicides and/or crop oils on broadleaf crops treated with Albaugh Rometri 480 SC Selective Herbicide may cause severe damage.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

Harmful if swallowed. Avoid inhaling spray mist. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or the Poisons Information Centre (telephone 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS) which is available from the supplier and accessible from the Albaugh website albaugh.com/au

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of strains resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but we cannot exercise any control over their mixing, use or application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application, or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

APVMA Approval Number: 69264/141466