

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ALBAUGH ZIPMAL 960 EC HERBICIDE

ACTIVE CONSTITUENT: 960 g/L S-METOLACHLOR

GROUP

15

HERBICIDE

Controls certain annual grasses and broadleaf weeds in certain crops, such as rhubarb, brassica leafy vegetables, mustard, spinach, silverbeet, spring onions, shallots, culinary herbs, as specified in the Directions for Use

IMPORTANT: READ THE ATTACHED LEAFLET/BOOKLET BEFORE USE

ALBAUGH AUSTRALIA PTY LTD

Level 1, 530 Little Collins Street,

Melbourne VIC 3000

Australia

Tel 03 9909 7183

www.albaugh.com/au

SELECTIVE

HERBICIDE

Manufactured globally to an American standard.

ALBAUGH

your alternative

DIRECTIONS FOR USE  
RESTRAINTS

DO NOT apply to waterlogged soils.  
DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 2 days of application.  
DO NOT irrigate to the point of runoff for at least 2 days after application.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)  
DO NOT allow bystanders to come into contact with the spray cloud.  
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.  
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.  
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Crop	Weeds Controlled	State	Rate	Critical Comments
Rhubarb	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	All states	1 - 2 L/ha	<b>Pre-emergent:</b> Apply one (1) application only, immediately before or after transplanting crowns and before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following re-hilling due to exposure of untreated soil.
<b>Brassica leafy vegetables</b> including Chinese white cabbage, Chinese flowering cabbage, Chinese broccoli, Indian mustard cabbage, Chinese cabbage, Mizuna greens, Red mustard, Curled mustard, Chinese flat cabbage, Mustard greens, Turnip greens, Rutabaga greens, and Mibuna greens)	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	All states	1 - 2 L/ha	Apply one treatment only, immediately after transplanting. Apply in a minimum of 60 L water per hectare. Apply sufficient irrigation after application to wet the upper 3 to 4 cm of soil with 24 hours. Use rates towards the higher end of the range where blackberry nightshade, chickweed, fat hen, love grass, pigweed, potato weed and wire weed are major problems (Refer Brassica use pattern on product label). See general instructions for warning – brassica crops.
Spinach and silverbeet	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	All states	1.5 – 2.0 L/ha	Apply with a boom spray, in a minimum of 60 L water/ha Apply immediately before, at, or immediately after sowing. If transplanting the crop, apply immediately before transplanting. DO NOT apply herbicide over the transplants. Crop retardation may occur where the herbicide is used on soils that have low organic matter content and which contain more than 60% of fine sand and silt. On these soil types, use rates towards the low end of the rate range, and apply to moist soil. Irrigation after application on these types of soils should be limited to no more than 25 mm. Also, on these soils, DO NOT use more than once per year (as on product label for brassicas); For other relevant instructions, see Broccoli (etc) section of Directions for Use table.
Spring onions ( <i>Allium fistulosum</i> ) and Shallots ( <i>Allium cepa</i> . var. <i>aggregatum</i> )	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	All states	1 L/ha	<b>DIRECT SEEDED CROPS:</b> Apply one application before, at or immediately after planting and before crops and weeds have germinated; <b>TRANSPLANTED CROPS:</b> Apply one application before planting and before weeds have germinated.

Crop	Weeds Controlled	State	Rate	Critical Comments
<b>Culinary herbs:</b> basil, bay leaves, borage, chives, coriander, dill, fennel, lemon balm, lemon grass, kaffir lime leaves, marigold flowers, marjoram (oregano), mints, nasturtium leaves, parsley, rosemary, sage, salad burnett, sorrel, tarragon, thyme, <b>savoury root vegetable:</b> galangal <b>leafy vegetables:</b> rucola (rocket), chervil, mizuna. <b>teas:</b> lemon verbena <b>spice:</b> turmeric. <b>edible flowers:</b> dianthus, roses.	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	All states	1.5 – 2.0 L/ha	Apply before, at, or immediately after planting/transplanting and before crops and weeds have germinated. Sufficient irrigation to wet the top 3-4 cm of soil should be applied within 24 hours. Do not use more than once per season.
Broccoli, Brussels sprouts, Cabbages, Cauliflowers	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)  Crowsfoot Grass, Fierce Thornapple (suppression), Liverseed Grass, Winter Grass  Apple of Peru, Awnless Barnyard Grass, Fierce Thornapple (suppression), Liverseed Grass, Yellow Weed, Redroot Amaranth (suppression).  Annual Ryegrass, Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Fiddle Dock, Italian Ryegrass, Liverseed Grass, Nettleleaf Goosefoot (suppression), Twigg Turnip (suppression), Winter Grass	NSW, ACT, Vic, Qld, Tas, NT only   NSW, ACT only  Qld only  Vic, Tas only	1.5 to 2 L/ha	Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Blackberry Nightshade, Chickweed, Fat-hen, Lovegrass, Pigeon Grass, Pigweed, Potato Weed and Wireweed are the major problem. See Warning - Brassica crops.  Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common Sowthistle, Fierce Thornapple, Liverseed Grass, Redroot Amaranth, Twigg Turnip and Nettleleaf Goosefoot are the major problems. <b>WARNING:</b> Qld. only stunting and reduced yield may occur in broccoli in red soils and light alluvial soils. See Warning - Brassica crops.
Canola and mustard (oilseed cultivars) ( <i>Brassica juncea</i> )	Toad Rush	NSW, ACT, SA, Vic, Tas, Qld and WA only	0.15 to 0.25 L/ha	Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil. Use rates towards the higher end of the range where longer residual control is required.
Cotton	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	NSW, ACT, Qld only	1 L/ha	<b>Pre-emergent:</b> Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas, as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where sand content of the soil is within the range 50% to 70%. See Warning - Cotton.
Green Beans, Navy Beans	Blackberry Nightshade, Common Sowthistle, Fat-hen, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass and suppression of Pigweed.  Powell's Amaranth	All states  Tas only	1.5 to 2 L/ha  2 L/ha	<b>Pre-emergent:</b> Apply before, at, or immediately after planting and before crops and weeds have germinated. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas, as this will reduce weed control. Do not use on soils where the sand content exceeds 70%. Use lower rate where the sand content is within the range 50% to 70%.  <b>Pre-emergent:</b> Apply before, at, or immediately after planting and before crops and weeds have germinated. For sufficient Amaranth control, the Albaugh ZIPMAL 960 EC Herbicide application should be followed by 1 L/ha Basagran* (2-leaf Amaranth) plus 2 L/ha Basagran* (4-leaf Amaranth). Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas, as this will reduce weed control.
Maize, Sweet Corn	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	NSW, ACT, Vic, Qld, SA, NT, Tas only	1 to 2 L/ha	<b>Pre-emergent:</b> Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas, as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range of 50% to 70%.
Pastures - Trikkala subclover, Haifa White clover, Maral Persian clover, Paradana Balansa clover, Redquin Red clover, Phalaris and Cocksfoot	Toad Rush	Vic, NSW, ACT, NT, SA, WA only	200 to 250 mL/ ha	Apply at or immediately after planting and before pasture species and weeds emerge. Application should be made to a moist seedbed. Sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days of spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required. DO NOT use in soils where prolonged waterlogging is anticipated, otherwise unacceptable crop retardation will occur. Avoid damage to pasture species by avoiding overlaps and double spraying in headlands where White, Persian and Balansa clovers are sown if the 250 mL rate is being used.

Crop	Weeds Controlled	State	Rate	Critical Comments
Peanuts	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Love Grass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, ACT, NT only	1 to 2 L/ha	Apply pre-irrigation where soil crusting is likely. Apply before, at or immediately after planting and before weeds germinate. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil.  If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use rate toward the higher end of the range on heavy soils or where a heavy grass population is expected.
Soybeans, Sunflowers				
Sorghum treated with oxabetrinil Sorghum Seed Safener	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	NSW, ACT, NT, Qld only	1 to 2 L/ha	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. DO NOT apply to sorghum seed not previously treated with oxabetrinil Sorghum Seed Safener. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable, incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.  Use rate toward the higher end of the range on heavy soils or where a heavy grass population is expected. Use the lower rate on sandy soils.
	Blackberry Nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1 to 2 L/ha plus 2.5 L/ha of atrazine (500 g/L) per ha	
Sugarcane plant and ratoon	Awnless Barnyard Grass, Barnyard Grass, Bellvine, Billygoat Weed (Blue Top), Blackberry Nightshade, Caltrop, Common Sowthistle (Milk Thistle), Common Thornapple, Crowsfoot Grass, Green Summer Grass, Guinea Grass, Mexican Clover, Mintweed, Needle Burr, Passionfruit Vines, Pigeon Grass, Pigweed, Redroot Amaranth, Square Weed, Star of Bethlehem, Summer Grass, Wild Rose	Southern Qld only (Bundaberg South)	1.1 to 1.45 L/ha plus 2.7 to 3.6 L/ha of an atrazine (500 g/L)	Apply as a pre- or post-emergent application to crop. Apply once per year only. Application should be made to a moist soil before weeds and grasses have germinated. If conditions remain dry for a period of 10 days after spraying, irrigation or a shallow cultivation (2.5 cm) may assist results. DO NOT throw untreated soil on to treated areas as this will reduce weed control. In Southern Queensland, use rates toward the higher end of the range on heavy soils where a high grass population is expected. In Northern Queensland, application must be made to moist soil and rainfall or irrigation should occur within 24 hours of application. Use rates toward the higher end of the range where high Green Summer Grass population is expected. Where broadleaf weeds and grasses have emerged and are in the 2 to 4 true leaf stage, knockdown herbicides such as ametryn (500 g/L) at 2 L/ha, flowable diuron (500 g/L) at 1 L/ha or paraquat (250 g/L) at 1.2 to 1.6 L/ha, applied as a post directed spray, can be added.
		Northern Qld only (Mackay North)	1.45 to 1.8 L/ha plus 3.6 to 4.5 L of an atrazine (500 g/L) or 2 to 2.5 kg/ha of atrazine (900 g/kg)	
Sweet Potatoes	Black pigweed, Red Shank, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrasses, Pigeon Grass, Summer Grass, Suppression of Evening Primrose	NSW, ACT, Qld, Vic, NT, SA only	1.5 L/ha	<b>Pre-emergent:</b> Apply immediately after transplanting before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours.  Further weed germination may occur following re-hilling due to exposure of untreated soil.
Tobacco (transplanted in field)	Barnyard Grass, Lovegrass, Pigeon Grass, Summer Grass and suppression of Common Thornapple ( <i>Datura stramonium</i> ) and Nightshades ( <i>Solanum sarrachoides</i> and <i>S. nigrum</i> )	Vic, NSW, ACT only	1.5 L/ha	Apply by low volume boom spray prior to transplanting and before weeds and grasses germinate. The seedbed should be fine and free of large clods. Thoroughly incorporate within 7 days of spraying using a rotary hoe or similar equipment set to work to a depth of 7.5 cm. DO NOT use disc-cultivating implements. An even coverage of the herbicide is essential. The herbicide needs to be activated by moisture through to the top 3 to 4cm of soil at the time of application or within 4 days of incorporation. This can be achieved by applying to moist soil, irrigating soon after incorporation or irrigating immediately after transplanting. Avoid boom overlaps as double treatment may cause growth suppression. Temporary crop retardation may be observed when soil conditions remain dry after application or where black root rot may occur.
Barley, Oats	Suppression of Annual Ryegrass	NSW, ACT, Vic, Tas, SA, WA only	375 to 500 mL/ ha	Apply to moist soils immediately after planting and before crop and weeds emerge. Use the lower rate on light soils. Application should be made to a moist seedbed and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Damage may occur if crop is sown too shallow (less than 4 cm).  Damage can occur on light sandy soil types after heavy rain. See General Instructions for use in zero-tillage systems.
Wheat, Barley, Oats, Triticale	Toad Rush	Vic, NSW, ACT, SA, WA only	150 to 250 mL/ ha	Apply at or immediately after planting and before crop and weeds emerge. Application should be made to a moist seedbed and sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required or in situations where crop competition is minimal.  When mixed with triasulfuron formulations, apply pre-sowing and incorporate by sowing operation (for wheat only).  See General Instructions for use in zero-tillage systems

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

SPINACH AND SILVERBEET:

GRAZING: DO NOT GRAZE OR USE FOR STOCKFEED FOR 8 WEEKS AFTER APPLICATION

CANOLA, MUSTARD: DO NOT GRAZE TREATED FIELDS FOR 10 WEEKS AFTER APPLICATION

GREEN BEANS, NAVY BEANS:

HARVEST: DO NOT HARVEST FOR AT LEAST 8 WEEKS AFTER APPLICATION

GRAZING: DO NOT GRAZE OR USE FOR STOCKFEED FOR AT LEAST 10 WEEKS AFTER APPLICATION

HERBS, CHERVIL, MIZUNA, RUCOLA, LEMON VERBENA: DO NOT HARVEST FOR AT LEAST 8 WEEKS AFTER APPLICATION

GALANGAL, TURMERIC: DO NOT HARVEST FOR AT LEAST 23 WEEKS AFTER APPLICATION

PASTURES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION

SWEET POTATOES: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 23 WEEKS AFTER APPLICATION

WHEAT, BARLEY, OATS, TRITICALE:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION

HARVEST withhold for all other crops

NOT REQUIRED WHEN USED AS DIRECTED

FORAGE withhold for all other crops (except Sweet Potatoes)

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION

Trade Advice:

Export of treated produce:

Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with this product. If you are growing edible produce for export, please check with ALBAUGH AUSTRALIA PTY LTD for the latest information on MRLs and import tolerances before using this product.

GENERAL INSTRUCTIONS

USES

Albaugh ZIPMAL 960 EC Herbicide is a short residual, pre-emergent herbicide for the control of many important annual grasses and broadleaf weeds in the crops nominated. The product does not control the important broadleaf weeds in maize, sweet corn and with oxabetrinil treated sorghum, but when mixed with atrazine (500 g/L) control of many important annual weeds and grasses is obtained without the problem of carryover of herbicides affecting succeeding crops. The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3 to 4 cm of soil, where it must be present at the time of weed germination.

APPLICATION

**Ground Application only:** Apply in a minimum of 60 L/ha of water. DO NOT apply by aircraft.

MIXING

Add the required amount directly to the spray tank, fill with water and mix well.

**Zero-tillage (Canola, Barley, Oats, Wheat, Triticale)**

When sowing using knife points and zero-tillage, and leaving a seeding slot, do not apply post-seeding unless harrows (or similar) have filled-in the seeding slot before application. Alternatively, apply Albaugh ZIPMAL 960 EC Herbicide immediately before seeding.

**Warning - Cotton:** Heavy rain following an application of Albaugh ZIPMAL 960 EC Herbicide, together with below average soil temperatures, can result in crop injury, due to the inability of the germinating plant being able to metabolise the herbicide. Albaugh ZIPMAL 960 EC Herbicide should not be used in situations where these conditions are likely to occur.

**Warning - Brassica Crops:** Crop retardation may occur where Albaugh ZIPMAL 960 EC Herbicide is used on soils that have a low organic matter content and which contain more than 60% of fine sand and silt. On these soil types, use rates of Albaugh ZIPMAL 960 EC Herbicide towards the lower end of the rate range and apply onto moist soil. Irrigation after application should be limited to no more than 25mm. DO NOT use more than once per year.

**To Avoid Crop Damage**

As metolachlor is known to cause some crop damage in sandy-sandy loam soils with low organic matter, it is recommended that growers test the safety of metolachlor on a small area of their crop before using the product on larger areas. If any adverse crop symptoms are observed within 14 days of application in which irrigation or rainfall has occurred, then the product should not be used.

RESISTANT WEEDS WARNING

GROUP	<b>15</b>	HERBICIDE
-------	-----------	-----------

Albaugh ZIPMAL 960 EC Herbicide is a member of the α-Chloroacetamides group of herbicides. Albaugh ZIPMAL 960 EC Herbicide has the inhibition of very long chain fatty acid synthesis (VLCFA inhibition) mode of action. For weed resistance management this is a Group 15 herbicide. Some naturally occurring weed biotypes resistant to Albaugh ZIPMAL 960 EC Herbicide and other Group 15 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Albaugh ZIPMAL 960 EC Herbicide or other Group 15 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, ALBAUGH AUSTRALIA PTY LTD accepts no liability for any losses that may result from the failure of Albaugh ZIPMAL 960 EC Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, or Primary Industries Department.

PRECAUTIONS

Re-entry Period

Do not allow entry into treated areas until the spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause drift onto nearby plants/crops, cropping lands or pastures.  
  
DO NOT plant crops other than broccoli, Brussels sprouts, cabbages, cauliflowers, cotton, maize, peanuts, soybeans, sunflowers, sweet corn, tobacco or oxabetrinil treated sorghum within 6 months of application of Albaugh ZIPMAL 960 EC Herbicide

PROTECTION OF LIVESTOCK

DO NOT graze or cut sweet potato leaves for stock food.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT TOXIC TO FISH.** DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

DO NOT apply under meteorological conditions, or from spraying equipment, which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, water bodies or watercourses lands or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well-ventilated area out of direct sunlight. Protect from frost.

Non-refillable containers

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply/designated collection point/other specific collection details for refill or storage.

SAFETY DIRECTIONS

Will damage the eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, and face shield or goggles, and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS) which is available from the supplier and accessible from the Albaugh website albaugh.com/au

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of strains resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but we cannot exercise any control over their mixing, use or application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application, or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

APVMA Approval Number: 94198/141806