

ACARAMIK® EC

**FOR THE CONTROL OF PEST MITES ON PEARS, APPLES,
TOMATOES, CITRUS, HOPS, STRAWBERRIES AND
ORNAMENTALS.**

**FOR THE CONTROL OF PEST MITES AND NATIVE BUDWORM
ON COTTON.**


After application, Albaugh ACARAMIK® EC Insecticide/Miticide quickly moves into leaves where it remains for several weeks and where it is taken up by sucking mites.

ACARAMIK® EC is not systemic making good coverage essential. Abamectin that is not absorbed into plants is quickly degraded.

PRODUCT OVERVIEW

- Do not make more than two applications to cotton per season, regardless of pest controlled.
- Apply by ground spraying equipment or fixed wing aircraft in a minimum volume of 20 litres per hectare.
- Good coverage is essential.
- Monitor crops regularly and apply as soon as the threshold mite number for your area has been reached.
- Best results will be obtained when applied to low mite populations.
- Application to high populations may not give satisfactory control. Under these conditions a second miticide application 7-10 days later may be needed.
- Do not use more than twice in one season for mite control.
- Refer to notes on resistance under General Instructions section of the ACARAMIK® EC label.
- Do not spray plants in flower while bees are foraging.

Sorcerer® is a registered trademark of Adama Australia



ACTIVE INGREDIENT

18 g/L Abamectin
266 g/L N-Methylpyrrolidone
60 g/L Hydrocarbon Liquid

I.R.A.C GROUP

GROUP 6 INSECTICIDE

FORMULATION

Emulsifiable Concentrate

POISON SCHEDULE

Schedule 6

COMPARE TO

Sorcerer®18

MODE OF ACTION

Glutamate-gated Chloride (GluCl)
channel allosteric modulators.
Nerve action.

SIGNAL HEADING

POISON
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

WATER RATES

Aircraft: 20L/ha of water
Horticultural applications use
1000L/ha with a horticultural oil

PACKAGING

5LTR, 10LTR

APVMA APPROVAL NO.

56346/141424

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,
WA