

BOGEN® 600 EC

**CONTROLS A VARIETY OF UNWANTED HARD TO KILL TREES,
BUSH AND VARIOUS BROADLEAF WEEDS.**



Albaugh BOGEN® 600 EC Herbicide is a member of the pyridyloxy-carboxylate group of herbicides. BOGEN® 600 EC has the disruptors of plant cell growth mode of action.

PRODUCT OVERVIEW

- **Application**
Apply with hand directed equipment such as high volume spraying equipment, knapsack sprayer, knapsack misting machines with the following exceptions:
Mistblower: Eucalyptus, Blackberry, Brigalow only
Aircraft: Blackberry only
Controlled Droplet Application (CDA): Blackberry only
Gas Gun, Sprinkler Sprayer: Eucalyptus seedlings only
Boomsprayer: Paddy Melon, Afghan, Bitter or Camel Melon only
- Highly effective on melon weed species.
- Weeds controlled include blackberry, camphor laural, English broom, gorse, common prickly pear, saffron thistle, capeweed and lantana (refer to the label to see full weed list).
- Fallow crop land weed control.
- Firebreak weed control.
- Commercial/industrial land weed control.

ACTIVE INGREDIENT

600 g/L Triclopyr

H.R.A.C. GROUP

GROUP 4 HERBICIDE

FORMULATION

Emulsifiable Concentrate

POISON SCHEDULE

Schedule 6

COMPARE TO

Garlon®

MODE OF ACTION

Synthetic auxin

SIGNAL HEADING

POISON
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

20LTR, 110LTR

APVMA APPROVAL NO.

94156/141713

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,
WA



Garlon® is a registered trademark of Corteva Agriscience