

COVERPRO® 480 SC

**A BROAD SPECTRUM, SYSTEMIC, FOLIAR APPLIED FUNGICIDE
FOR THE CONTROL OF VARIOUS DISEASES IN CANOLA AND
CEREAL CROPS**

Up to two sprays of Albaugh COVERPRO® 480 SC may be applied per season to the crop.

DISEASES CONTROLLED

- Blackleg
- Net form Net Blotch (*Pyrenophora teres f. teres*)
- Spot form Net Blotch (*Pyrenophora teres f. maculata*)
- Strip Rust (*Puccinia striiformis*)
- Sclerotinia - generally a single application of Albaugh COVERPRO® 480 SC at 20 to 30% flowering will control sclerotinia in crops with a short flowering interval. Crops with an extended flowering period may require a second application prior to 50% flowering (full-bloom) to adequately control sclerotinia if conditions late in the season are conducive to development of disease.
- Yellow Leaf Spot (*Pyrenophora tritici-repentis*)
- Septoria Nodorum-glume Blotch (*Phaeosphaeria nodorum*)
- Leaf Rust (*Puccinia coronata f. sp. avenae*)
- Septoria Blotch (*Phaeosphaeria avenaria*)
- Use with a non-ionic surfactant at 0.25% v/v



ACTIVE INGREDIENT

480 g/L Prothioconazole

F.R.A.C. GROUP

GROUP 3 FUNGICIDE

FORMULATION

Suspension Concentrate

POISON SCHEDULE

Schedule 5

COMPARE TO

Various

MODE OF ACTION

DMI Inhibitor

SIGNAL HEADING

CAUTION
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

20LTR, 110LTR, 1000LTR

APVMA APPROVAL NO.

95990/147834

STATES AND TERRITORIES REGISTERED

NSW, NT, QLD, SA, TAS, VIC, WA