

DITTO[®]

CONTROLS WILD OATS, PARADOXA GRASS (ANNUAL PHALARIS), CANARY GRASS AND ANNUAL RYEGRASS IN WHEAT. HIGHLY ACTIVE ON WILD OATS AND CAN BE MIXED WITH MANY BROADLEAF HERBICIDES.



PRODUCT OVERVIEW

- Apply to wheat at 2 leaf to late jointing (Zadocks 12-37).
- May be tank mixed with compatible herbicides.

KEY WEEDS CONTROLLED OR SUPPRESSED

Wild oats

Paradoxa grass (*Phalaris paradoxa*)

Canary grass (*Phalaris minor*)

Annual ryegrass

Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.

KEY CROPS

Wheat

ACTIVE INGREDIENT

240 g/L Clodinafop-propargyl
60 g/L Cloquintocet-mexyl

H.R.A.C. GROUP

GROUP 1 HERBICIDE

FORMULATION

Emulsifiable Concentrate

POISON SCHEDULE

Schedule 6

COMPARE TO

Topik[®]

MODE OF ACTION

Inhibitors of acetyl CoA
carboxylase

SIGNAL HEADING

POISON
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

5LTR

APVMA APPROVAL NO.

81130/141541

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,
WA



Topik[®] is a registered trademark of Syngenta