

FLUROPIX® 400 EC

A SELECTIVE HERBICIDE FOR POST-EMERGENT BROADLEAF
WEED CONTROL



PRODUCT OVERVIEW

- Use Albaugh FLUROPIX® 400 EC Herbicide for the control of a wide range of broadleaf weeds in fallow, lucerne, maize, millets, pastures, poppies, sorghum, sugar cane, sweet corn, winter cereals. Also for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, forests, pastures and rights-of-way.
- Controls annual broadleaf weeds.
- FLUROPIX® 400 EC is a systemic herbicide that moves within the plant, controlling exposed and underground plant tissue.
- Broad tank mix flexibility with most grass and broadleaf herbicides.

KEY WEEDS CONTROLLED OR SUPPRESSED

Bathurst burr, noogoora burr, black bindweed (climbing buckwheat), mimosa pigra, bellyache bush, blackberry nightshade, bokhara clover, caltrop (yellow vine) (tribulus terrestris) (T. micrococcus), cobblers pegs, cockspur thorn, creeping lantana, crofton weed, mistflower, docks (rumex spp.), hexham scent, honey locust, small flowered mallow (marshmallow) (malva parviflora), lantana.

Woody weed situations: Weeds must be actively growing to attain optimal effect. Delay the treatment of regrowth following bulldozing, slashing, burning, ploughing or a previous chemical treatment until it has at least 1 metre of new, vigorous, growth.



See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

400g/L Fluroxypyr present as the Methyl Heptyl Ester

H.R.A.C. GROUP

GROUP 4 HERBICIDE

FORMULATION

Emulsifiable Concentrate

POISON SCHEDULE

Schedule 5

COMPARE TO

Various

MODE OF ACTION

Pyridine group, disrupters of plant cell growth

SIGNAL HEADING

CAUTION
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

20LTR, 110LTR, 1000LTR

APVMA APPROVAL NO.

81092/141581

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA