

HEXANIL[®] 250 SL

**FOR THE CONTROL OF CERTAIN BROADLEAF WEEDS,
PERENNIAL AND ANNUAL GRASSES, WOODY WEEDS IN PINUS
RADIATA PLANTATIONS, PASTURE SITUATIONS AND
COMMERCIAL AND INDUSTRIAL AREAS AND RIGHTS OF WAY.**



PRODUCT OVERVIEW

- Albaugh HEXANIL[®] 250 SL Herbicide is both foliar and root absorbed and will give residual soil activity. For best foliar and residual control, apply during periods of active plant growth.
- Application when vegetation is dormant, semi-dormant, under moisture stress or during winter months may not be as effective.
- In temperate winter rainfall areas where temperatures are generally below 25°C during periods of active growth, overall effect may be slower to appear.
- For best control of perennials, apply during the months of spring in these areas.
- In WA and SA apply only during April-August when adequate rainfall is expected.
- Sufficient moisture is required after treatment to promote root absorption by the weed species.

KEY USES

Pinus radiata.

Commercial and industrial areas, rights of way, around agricultural buildings and pasture situations.

ACTIVE INGREDIENT

250 g/L Hexazinone

H.R.A.C. GROUP

GROUP 5 HERBICIDE

FORMULATION

Soluble Concentrate

POISON SCHEDULE

Schedule 5

COMPARE TO

various

MODE OF ACTION

Triazinones; inhibitors of photosynthesis at photosystem II

SIGNAL HEADING

CAUTION
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

10LTR, 20LTR

APVMA APPROVAL NO.

81083/141683

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,
WA



See label for complete application guidelines and restrictions.