RUMBA® 750 SG

FOR THE CONTROL OF A WIDE RANGE OF BROADLEAF WEEDS IN WHEAT, BARLEY, OATS, TRITICALE, CANOLA, PASTURES AND FALLOW LAND.



PRODUCT OVERVIEW

- Strong tool for control of California thistle and other broadleaf weeds in many key crops.
- Trusted formulation with excellent crop safety.
- Many tank-mix partner options to tailor your application needs.

KEY WEEDS CONTROLLED OR SUPPRESSED

Plantback periods NSW, Vic, SA, WA (winter rainfall areas)

Albaugh RUMBA® 750 SG Herbicide Rate g/ha	Up to 120	200	>200
Chickpea, field pea, faba bean, lupins, medics & clover	9 months	12 months	24 months
Wheat, barley, oats	1 week	-	-

Plantback period NSW, Qld (summer rainfall areas)

Albaugh RUMBA® 750 SG Herbicide Rate g/ha	30	60	120
Wheat, barley, oats	1 week	1 week	
Chickpea		12 weeks	
Lucerne	36 weeks	36 weeks	36 weeks
Cotton	2 weeks	4 weeks	8 weeks
Sorghum, maize	1 week	2 weeks	2 weeks
Sunflower	5 weeks	8 weeks	24 weeks
Soybean	1 week	1 week	24 weeks

For plantback periods of >4 weeks, 100mm rain must have fallen between application of RUMBA® 750 SG and planting susceptible crop.



ACTIVE INGREDIENT

750 g/kg Clopyralid present as the Potassium Salt

H.R.A.C. GROUP

GROUP 4 HERBICIDE

FORMULATION

Water Soluble Granules

POISON SCHEDULE

Schedule 5

COMPARE TO

Lontrel®

MODE OF ACTION

Pyridines: disrupters of plant cell growth

SIGNAL HEADING

CAUTION KEEP OUT OF REACH OF **CHILDREN READ SAFETY DIRECTIONS** BEFORE OPENING OR USING

PACKAGING

2KG

APVMA APPROVAL NO.

94650/143090

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA



