

TEBIX[®] 200 GR

**HIGHLY CONCENTRATED HERBICIDE FOR EXCELLENT
BROADLEAF WEED CONTROL**



PRODUCT OVERVIEW

- Use Albaugh TEBIX[®] 200 GR Herbicide for control of brigalow regrowth, tea tree regrowth, mimosa pigra and certain problem woody weeds on grazing lands by hand, aerial and ground application as specified.

KEY USES

Use of TEBIX[®] 200 GR on native vegetation must be done in accordance with State and/or Local legislation.

Protection of livestock: Poisonous plants may become more palatable after application of granules and stock should be kept away from these plants until they have died down.

TEBIX[®] 200 GR may be applied at any time of the year. Treatments become effective after sufficient rainfall has occurred to move the chemical into the root zone where it is absorbed by woody plants. Applications just prior to seasonal rainfall give the most rapid response.

- Herbicidal activity will continue for several seasons after application. During this time woody plants may go through repeated defoliations and regrowths until death occurs. A single application is normally effective for several years.
- Forage grass production usually increases as woody weed competition is reduced. Increased grass production is also dependent on adequate rainfall and an acceptable grazing management programme.
- It is important to allow grasses to seed and seedlings to become established after treatment before grazing pressure is increased. Areas treated with TEBIX[®] 200 GR may be overseeded with suitable pasture species.
- Consult your agricultural specialist for details on suitable seeding rates, timing and fertiliser application programme.
- The residual action of TEBIX[®] 200 GR may be impaired if fire occurs before clay pellets are dissolved and the chemical has moved into the root zone.

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

200 g/kg Tebuthiuron

H.R.A.C. GROUP

GROUP 5 HERBICIDE

FORMULATION

Granular Formulation

POISON SCHEDULE

Schedule 6

COMPARE TO

Various

MODE OF ACTION

Ureas group of herbicides.
Inhibitors of photosynthesis at
photosystem II

SIGNAL HEADING

POISON
KEEP OUT OF REACH OF
CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING

PACKAGING

5KG, 20KG, 1000KG

APVMA APPROVAL NO.

94747/143525

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,
WA