

TUKEN[®] EXTRA

FOR THE CONTROL OF CERTAIN BROADLEAF WEEDS IN WINTER CEREALS AND CLOVER AS SPECIFIED



PRODUCT OVERVIEW

- Cereals: Wheat, barley, oats, triticale, cereal rye.
- Pasture: Newly sown and established clover-based pasture, clover for hay and seed production.
- Cereals: After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.
- Oats: The tolerance of varieties Esk and Nile (the two main varieties grown in Tasmania) to Albaugh TUKEN[®] Extra Herbicide has not been tested. Test a small area of crop before using TUKEN[®] Extra over large areas. Consult your local Albaugh representative for advice on specific varieties.

KEY WEEDS CONTROLLED

Canola (Rape), Brassica napus
 Charlock, Sinapis arvensis
 Common Sowthistle (Milk Thistle), Sonchus oleraceus
 Cowvine, Ipomoea lonchophylla
 Deadnettle, Lamium amplexicaule
 Dock, Rumex spp.
 Fat Hen, Chenopodium album
 Fumitory, Fumitory spp.
 Hexham Scent (King Island Melilot), Melilotus indicus
 Hyssop Loosestrife, Lythrum hyssopifolia
 Indian Hedge Mustard, Sisymbrium orientale
 Long Storksbill, Erodium botrys
 Mouse-eared Chickweed, Cerastium glomeratum
 Paterson's Curse, Echium plantagineum
 Prickly Lettuce, Lactuca serriola
 Rough Poppy, Papaver hybridum
 Scarlet Pimpernel, Anagallis arvensis
 Skeleton Weed, Chondrilla juncea
 Stemless Thistle, Onopordum acaulon
 Tree Hogweed, Polygonum patulum
 Variegated Thistle, Silybum marianum
 Volunteer Lupins, Lupinus spp.
 Wild Radish, Raphanus raphanistrum
 Wireweed (Hogweed), Polygonum aviculare

Capeweed, Arctotheca calendula
 Chickweed, Stellaria media
 Corn Gromwell, Buglossoides arvensis
 Crassula, Crassula spp.
 Dense-flower Fumitory, Fumaria densiflora
 Doublegee (Spiny Emex), Emex australis
 Fireweed, Senecio spp.
 Hedge Mustard, Sisymbrium officinale
 Horehound, Marrubium vulgare
 Iceplant, Mesembryanthemum spp.
 London Rocket, Sisymbrium irio
 Marshmallow, Malva parviflora
 Night-scented Stock, Matthiola longipetala
 Peppergrass, Lepidium spp.
 Purple Goosefoot, Scleroblitum atriplicinum
 Saffron Thistle, Carthamus lanatus
 Shepherd's Purse, Capsella bursa-pastoris
 Sorrel, Rumex spp.
 Toad Rush, Juncus bufonius
 Turnip Weed, Rapistrum rugosum
 Vetch (Tares), Vicia sativa
 Ward's Weed, Carrichtera annua
 Wild Turnip, Brassica tournefortii



Jaguar[®] is a registered trademark of Bayer

ACTIVE INGREDIENT

250 g/L MCPA present as the Ethyl Hexyl Ester
 25 g/L Diflufenican

H.R.A.C. GROUP

GROUP 4 12 HERBICIDE

FORMULATION

Emulsifiable Concentrate

POISON SCHEDULE

Schedule 5

COMPARE TO

Jaguar[®]

MODE OF ACTION

Phenoxy and nicotinanalide groups, acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth

SIGNAL HEADING

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PACKAGING

5LTR, 10LTR, 20LTR, 110LTR, 1000LTR

APVMA APPROVAL NO.

81044/141684

STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA