



**NON-SELECTIVE  
HERBICIDE**

# ZIPPER® 200 EC

**FOR NON-RESIDUAL CONTROL OF BROADLEAF AND GRASS  
WEEDS IN VARIOUS SITUATIONS AS SPECIFIED**



- Albaugh ZIPPER® 200 EC Herbicide is a non-volatile herbicide with non-selective activity against many annual and perennial broadleaf weeds and grasses.
- ZIPPER® 200 EC is absorbed by plant foliage and green stems. It is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray.
- ZIPPER® 200 EC does not provide residual weed control. Visible symptoms of control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions.
- Best results are achieved when application is made under good growing conditions.
- Application to weeds under stress (e.g. due to continuous severe frosts, dry or waterlogged conditions) should be avoided.

## PRODUCT OVERVIEW

- **Compatibility:** ZIPPER® 200 EC is compatible with most residual herbicides, e.g. simazine, diuron, oxyfluorfen (Goal®), norfluazuron (Solicam®) and oryzalin (Surflan®). The addition of a wetting agent or other adjuvant is generally not considered necessary. However, benefit has been obtained using a wetting agent on hard-to-wet weeds when using water rates in excess of 500 L/ha. The rate is 25 mL/100 L of a 1000 g/L non-ionic wetting agent, or equivalent.

Liberty® is a registered trademark of BASF



### ACTIVE INGREDIENT

200 g/L Glufosinate-Ammonium

### H.R.A.C. GROUP

GROUP 10 HERBICIDE

### FORMULATION

Emulsifiable Concentrate

### POISON SCHEDULE

Schedule 5

### COMPARE TO

Liberty®

### MODE OF ACTION

Glycine group inhibitor of glutamine synthetase

### SIGNAL HEADING

CAUTION  
KEEP OUT OF REACH OF  
CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING

### PACKAGING

20LTR, 1000LTR

### APVMA APPROVAL NO.

81839/141371

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,  
WA