FUNGICIDE

PROPISTAR®

IMPROVING CROP PERFORMANCE WITH EARLY SEASON LEAF DISEASE PROTECTION.



PRODUCT FEATURES

- Wide window of application from cereal herbicide timing to full flag leaf.
- Early season control of blackleg in canola.
- Broad-spectrum disease control over a wide range of crops.

APPLICATION INFORMATION

Rate: 60 to 120 mL/ac

Water volume: Ground application: Minimum 80 L/ac (20 gal/ac)
Aerial application: 20 L/ac (5 gal/ac)

Acres treated: 40 to 80 acres/jug
Canola: 40 acres/jug

Wheat: 80 acres/jug for Septoria, 40 acres/jug for rust, leaf/tan spot
Oats: 40 acres/jug
Soybean/dry bean: 40 acres/jug

Surfactant: Not required

Rainfastness: 1 hour

TANK MIXABILITY

Herbicide tank-mixing – wheat & barley: PropiStar can be tank-mixed with only one of these herbicides at a time: 2,4-D Amine or Ester, Badge®, MCPA Amine or Ester, or Bromotril® 240 EC. See tank-mix partner label for rates to use.

Tank-mixing precautions:

- **Do not tank-mix PropiStar with herbicides for application onto oats.**
- Weeds and crops must be at the correct stage of growth as specified on both the PropiStar label and the tank-mix partner label.
- 2,4-D and MCPA formulations may be applied either by ground application or aerial application; tank mixes of PropiStar and Badge® or Bromotril® 240 EC can only be applied by ground application.
- Consult the herbicide partner label for a list of weeds controlled, directions for use, and precautions.
- Compatibility should always be confirmed by premixing small proportional quantities of water, PropiStar, and the tank-mix partner in advance.

Refer to label for full list of diseases controlled, complete application guidelines, and restrictions.

KEY CROPS
Wheat, barley, oats, canola, corn, soybeans, dry edible beans

ACTIVE INGREDIENT
Propiconazole

FUNGICIDE GROUP
3

FORMULATION
432 g/L EC

COMPARE TO
Bumper®, Tilt®

MODE OF ACTION
Sterol biosynthesis inhibitor

PACKAGING
4 x 4.8 L

REGISTRATION NUMBER
35564

280 Midpark Way SE #112 | Calgary, AB T2X 1J6 | albaugh.com/ca

ALBAUGH® your alternative



PROPISTAR®

MIXING INSTRUCTIONS

Herbicide mixing and spraying instructions:

- Spray equipment should be thoroughly flushed with clean water before mixing PropiStar.
- Fill spray tank 1/2 full with clean water. Engage gentle agitation.
- Add the required amount of PropiStar and agitate thoroughly.
- Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tank mix partner.
- Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.
- Use nozzle screens no finer than 50 mesh. Keep the by-pass line on or near the bottom of the tank to minimize foaming.

Fertilizer mixing and spraying instructions:

- If desired, small amounts of nitrogen may be applied with PropiStar.
- The appropriate amount of urea can be dissolved in water and added to the spray tank before adding PropiStar. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing PropiStar with a herbicide.

NOTE: Do not graze livestock on treated crops.

KEY DISEASES CONTROLLED OR SUPPRESSED*

Blackleg	Northern leaf blight	Spot blotch
Crown rust	Powdery mildew	Stem rust
Frogeye leaf spot	Scald	Tan spot
Leaf rust	Septoria glume blotch	
Net blotch	Septoria leaf blotch	

**Check label for specific crops and rates.*

◀ Refer to diseases for full list of diseases controlled, complete application guidelines, and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh, the A logo, and PropiStar are registered trademarks of Albaugh, LLC. Badge, Bromotril, and Bumper are registered trademarks of Adama. Tilt is a registered trademark of Syngenta. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. It is a violation of federal law to use any pesticide product other than in accordance with its labeling.

280 Midpark Way SE #112 | Calgary, AB T2X 1J6 | albaugh.com/ca

ALBAUGH® your alternative