

WARLOCK®

IMPORTANT COMPONENT OF INTEGRATED VEGETATION MANAGEMENT PROGRAMS DESIGNED TO RENOVATE OR RESTORE DESIRED PLANTS.



PRODUCT FEATURES

- Provides pre- and post-emergence control of emerging seedlings of susceptible weeds and regrowth of certain perennial weeds following application.
- Provides control of susceptible weeds when applied according to labeled directions.
- Compatible with certain labeled tank mixes with other brush-control herbicides.

APPLICATION INFORMATION

Rate: 0.10 to 0.20 L/ac

Water volume: Ground application:

Minimum 40 L/ac (10 gal/ac)

Aerial application: 7.7 L/ac (2 gal/ac)

Acres treated: 15 to 30 acres/jug

Surfactant: Consult label for specific tank mixes.

Rainfastness: Avoid application when heavy rain is forecast

TANK MIXABILITY

Warlock Plus 2,4-D Amine: Warlock can be tank mixed with 2,4-D Amine herbicide to broaden the spectrum of weeds controlled.

Warlock Plus Plotter®: Apply when targeted undesirable vegetation is actively growing. Only weeds and brush present at the time of application will be controlled.

Warlock can be tank mixed with either Arsenal® herbicide or glyphosate herbicides, present as isopropylamine salt, diammonium salt, potassium salt, or dimethylamine salt and registered for use on non-cropland areas, to broaden the spectrum of broadleaf weeds and invasive plants controlled.

MIXING INSTRUCTIONS

To prepare the spray mix, add half the desired amount of water to the spray tank. Then, with agitation, add the recommended amount of Warlock, followed by the tank-mix partner. If required, add the recommended amount of surfactant. Finally, with continued agitation, add the remaining water.

KEY CROPS

Rangeland, permanent pasture, industrial, and other non-crop areas

ACTIVE INGREDIENT

Aminopyralid

HERBICIDE GROUP

4

FORMULATION

SC

COMPARE TO

Milestone®

MODE OF ACTION

Synthetic auxin

PACKAGING

4 X 3.1 L

REGISTRATION NUMBER

34903

WARLOCK®

PRECAUTIONS

GENERAL USE PRECAUTIONS: Warlock is highly active against broadleaf plants. This product is recommended for use on areas where loss of broadleaf forage plants, including legumes, can be tolerated. Do not apply more than 0.2 L/ac of Warlock per annual growing season in rangeland, permanent pasture, industrial, and other **non-crop areas**.

SPRAYER CLEAN-OUT INSTRUCTIONS DO NOT use spray equipment used to apply Warlock for other applications to land planted to, or to be planted to susceptible crops or desirable sensitive plants, unless it has been determined that all residue of this herbicide has been removed by thorough cleaning of equipment. Equipment used to apply Warlock should be thoroughly cleaned before reusing to apply any other chemicals.

KEY WEEDS CONTROLLED

Absinth (wormwood)	Cudweed	Musk or nodding thistle	Skeletonweed
Bitter sneezeweed	Curly dock	Orange hawkweed	Spotted knapweed
Bull thistle	Dandelion	Oxeye daisy	Sulphur cinquefoil
Canada fleabane	Diffuse knapweed	Perennial sowthistle	Tall buttercup
Canada goldenrod	Fireweed	Plumeless thistle	Tall ironweed
Canada thistle	Fuller's teasel	Prickly lettuce	Tansy ragwort
Clover	Hairy buttercup	Purple loosestrife	Tropic croton
Common broomweed	Hairy fleabane	Rush	Tropical soda apple
Common ragweed	Horsenettle	Russian knapweed	Western ragweed
Common tansy	Japanese knotweed	Scentless chamomile	Yellow star thistle
Common yarrow	Mullein	Scotch thistle	



◀ Refer to label for full list of weeds controlled, complete application guidelines, and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh, the A logo, Plotter, and Warlock are registered trademarks of Albaugh, LLC. Milestone is a registered trademark of Corteva. Arsenal is a registered trademark of BASF. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. It is a violation of federal law to use any pesticide product other than in accordance with its labeling.