

# EVENTEX™ 400SC

**INDUSTRY STANDARD ACTIVE INGREDIENT  
FOR RESIDUAL CONTROL OF LEPIDOPTERAN  
INSECT PESTS.**

Eventex™ 400SC insecticide is a member of the anthranilic diamide class of insecticides with a mode of action acting on insect ryanodine receptors. Although Exceliprole has contact activity, it is most effective through ingestion of treated plant material. After exposure to Eventex, affected insects will rapidly stop feeding, become paralyzed, and typically die within 1-3 days. Time applications to the most susceptible insect pest stage, typically at egg lay, egg hatch and/or newly hatched larvae, before populations reach damaging levels. If possible, make applications at or before egg deposition to be most effective in minimizing damage levels caused by insect pests. When pest populations are high, use the highest listed application rate for that pest.

## PRODUCT FEATURES

- Active against a broad-spectrum of Lepidopteran pests, including navel orangeworm, peach twig borer, codling moth, and Oriental fruit moth.
- Fast-acting control, with feeding cessation occurring minutes after exposure.
- Extended residual control.
- Single active ingredient with minimal PPE requirements, short REI and short PHI for maximum flexibility.
- Minimal impact on beneficial insects, when used in accordance with sound integrated pest management practices.

See label for complete application guidelines and restrictions.



### ACTIVE INGREDIENT

3.34 lbs/gal chlorantraniliprole

### IRAC GROUP

28

### FORMULATION

Suspension concentrate

### REI

4 hrs

### COMPARE TO

Shenzi® 700WG

### MODE OF ACTION

Ryanodine receptor modulator

### SIGNAL WORD

Caution

### RESTRICTED USE

No

### PACKAGING

4 x 1 gal

### EPA REG NUMBER

83100-77 | AD110624

# EVENTEX™ 400SC

## FREQUENTLY ASKED QUESTIONS

### How should Eventex™ 400SC be applied to pome fruits for best results?

- Apply 2.1-3.75 fl oz/A by foliar application for control of labeled pests. Refer to label for state restrictions regarding pests and/or specific crops.
- Thorough coverage is essential. Select a spray volume appropriate for the size of trees or plants and density of foliage. For ground applications, apply in 30 or more gallons of water per acre, but DO NOT exceed 200 gallons per acre. For best results apply 100 – 150 gal water per acre.
- PHI: 5 days
- Apply higher rates within the listed range for heavier infestations, larger/denser crops or extreme environmental conditions such as rainy weather and high temperatures.
- Do not make more than three applications per calendar year. Minimum interval between applications is 10 days.

### How should Eventex™ 400SC be applied to stone fruits for best results?

- Apply 2.5-3.75 fl oz/A by foliar application for control or suppression of labeled pests.
- Thorough coverage is essential. Select a spray volume appropriate for the size of trees or plants and density of foliage. For ground applications, apply in 30 or more gallons of water per acre, but DO NOT exceed 200 gallons per acre. For best results apply 100 – 150 gal water per acre.
- PHI: 10 days
- Apply higher rates within the listed range for heavier infestations, larger/denser crops or extreme environmental conditions such as rainy weather and high temperatures.
- Do not make more than four applications per acre per crop, or eight applications per acre per calendar year. Minimum interval between applications is seven days.



◀ See label for complete application guidelines and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks and Eventex is a trademark of Albaugh, LLC. Shenzi is a trademark of UPL Ltd. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.