

INDOLEX™ 1.25EC

**FAST-ACTING, LONG-LASTING, BROAD-SPECTRUM
CONTROL OF A RANGE OF LEPIDOPTERAN INSECTS
AND OTHER KEY PESTS.**



Indolex™ 1.25 EC insecticide is an emulsifiable concentrate that can be applied as a foliar spray to control many important Lepidopteran and other key insects in a range of crops.

Apply Indolex 1.25 EC at the rates listed on the label when insect populations reach locally determined economic thresholds. Applications should target eggs and small instar larvae. Follow-up treatments should be applied, as needed, to keep pest populations within threshold limits. Apply at 5- to 7-day intervals or as needed to manage specific target insect pests, as specified in the specific crop sections of the product label, to maintain control.

PRODUCT FEATURES

- Active against a range of Lepidopteran insects, as well as corn rootworm adults, alfalfa weevil larvae, potato leafhopper, armyworm species, and grasshoppers
- Provides fast knockdown by contact and ingestion, causing irreversible feeding cessation.
- Provides fast knockdown with residual control.
- Helps conserve certain beneficial insects, including parasitic wasps, predatory mites, big-eyed bugs, damsel bugs, minute pirate bugs, and spiders.
- Short pre-harvest interval of 1-21 days, depending on crop.
- Excellent fit in integrated pest management and resistance management programs.

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

1.25 lbs/gal indoxacarb

IRAC GROUP

22A

FORMULATION

Emulsifiable concentrate

REI

12 hours

COMPARE TO

Steward® EC

MODE OF ACTION

Voltage-dependent sodium channel blockers (oxadiazine)

SIGNAL WORD

Warning

RESTRICTED USE

No

PACKAGING

4 x 1 gal, 2 x 2.5 gal

EPA REG NUMBER

83100-83 | AD021925

INDOLEX™ 1.25 EC

FREQUENTLY ASKED QUESTIONS

How should Indolex 1.25 EC be used on alfalfa for best results?

- For grasshopper and cabbage looper (except California) control, apply 4.6-11.3 fl oz/A.
- For alfalfa caterpillar, alfalfa weevil larvae, beet armyworm, cabbage looper (California only), Egyptian alfalfa weevil larvae, and granulate cutworm control, apply 6.7-11.3 fl oz/A.
- For potato leafhopper (except California) and western yellowstriped armyworm control, and Lygus bug suppression (Western U.S.), apply 9.2-11.3 fl oz/A.
- Apply lower listed rates for light to moderate infestations. Use intermediate to high rates within the listed rate range on heavier infestations or when later instar larvae exist. Use the highest listed rate for controlling severe infestations or when longer residual control is desired.
- Preharvest interval (defined as when the alfalfa crop is cut): 7 days
- Refer to label for additional use restrictions.

How should Indolex 1.25 EC be used on corn for best results?

- For beet armyworm, corn earworm, corn rootworm adult, European corn borer, fall armyworm, grasshopper, western bean cutworm, and yellowstriped armyworm control and Japanese beetle adult suppression, apply 6.0-11.3 fl oz/A.
 - Corn earworm control is only for treated foliage and silks. New foliage and new silks will not be protected with a single application.
- For brown stink bug, green stink bug, and southern green stink bug suppression, apply 9.2-11.3 fl oz/A.
- Preharvest interval: 14 days for grain and stover (field, pop, and seed corn) and 1 day for forage, fodder, and silage (field and seed corn only)
- Refer to label for additional use restrictions.



◀ See label for complete application guidelines and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks and Indolex is a trademark of Albaugh, LLC. Steward is a registered trademark of FMC Corporation. Always read and follow label restrictions, precautions, and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.