

MITICIDE/INSECTICIDE

ABAMECTIN 0.15 SELECT™

FOR THE CONTROL OF MITES, LEAF MINERS,
AND OTHER INSECTS.

Abamectin 0.15 Select™ ornamental miticide/insecticide is used to control mites and leaf miners on shadehouse, greenhouse, and field grown ornamentals. **Abamectin 0.15 Select** provides an excellent option for nursery professionals needing options for insecticide resistance management.

PRODUCT FEATURES

- Highly effective insecticide used to control mites and leafminers, among other pests.
- Safe to mix with horticultural spray oils for improved performance on certain species.
- Effective tool for combating pyrethroid-resistant insects.

LABELED USES

- Shadehouse, greenhouse, and field-grown ornamentals
- Christmas trees
- Woody ornamentals

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

2.0% Abamectin

IRAC GROUP

6

FORMULATION

Liquid concentrate

CHEMICAL FAMILY

Avermectin

COMPARE TO

Avid®

MODE OF ACTION

Inhibits nerve and muscle cell communication, paralyzing insect

RESTRICTED USE

No

SIGNAL WORD

Warning

PACKAGING

12 x 1 qt, 4 x 1 gal

EPA REG NUMBER

83100-5-89442

ABAMECTIN 0.15 SELECT™

FOR THE CONTROL OF MITES, LEAF MINERS, AND OTHER INSECTS

APPLICATION INFORMATION

Foliar Application

Abamectin 0.15 Select™ is an emulsifiable concentrate which when mixed with water according to the directions for use will control leafminers and mites and suppress whiteflies, thrips and aphids on ornamentals.

Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. For mites, apply when mites first appear and repeat as necessary to maintain control. For leafminers, apply as needed and repeat at 7-day intervals or as necessary to maintain control.

For suppression of aphids, thrips and whiteflies, apply when young, immature stages of these pests are first observed and repeat every 7 days for two or three weeks. After which time, rotate to other products that have different modes of action than **Abamectin 0.15 Select** for at least two to three weeks.

FOLIAR APPLICATION RATES AND TIMING

Shadehouse, greenhouse, field-grown ornamentals, foliage plants, Christmas trees, and other woody ornamentals		
PESTS	fl oz Abamectin 0.15 Select / 100 gals	COMMENTS
MITES: Carmine spider mite European red mite Southern red mite Spruce spider mite Two-spotted spider mite TARSONEMID MITES: Broad mites and Cyclamen mites ERIOPHYID MITES: Bud mites and Rust mites	4	For Tarsonemid mites, repeat applications to newly developing tissue may be necessary to maintain control.
<i>Liriomyza</i> leafminers	8	Repeat at 7-day intervals or as necessary to maintain control.
Boxwood leafminer	8	For control of mining larvae, make the application when adults are beginning to lay eggs in the new foliage
Aphids, thrips, and whiteflies	8	For suppression of pest populations, young immatures must be contacted by the spray.

Refer to label for full application and rate information.

LABELED MITES

Broad mite	Cyclamen mite	Southern red mite
Bud mites	European red mite	Spruce spider mite
Carmine spider mite	Rust mites	Two-spotted spider mite

LABELED INSECTS

Aphids	Thrips
Boxwood leafminer	Whiteflies
<i>Liriomyza</i> leafminer	



◀ See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. All rights reserved. Albaugh and the A logo are registered trademarks and Abamectin 0.15 Select is a trademark of Albaugh, LLC. Avid is a registered trademark of Syngenta. Always read and follow label restrictions, precautions, and directions for use. Products may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.