

FUNGICIDE

OXEM™ 4 ME SELECT

SYSTEMIC, PREVENTIVE, AND CURATIVE FUNGICIDE
FOR TURF AND ORNAMENTALS.

Oxem™ 4 ME Select is a systemic turf and ornamental fungicide containing mefenoxam for the control of damping off, root, and stem diseases caused by *Pythium* and *Phytophthora* spp., and foliar diseases such as downy mildew and those caused by *Phytophthora* spp. It works by suppressing essential components of the disease so the infection doesn't grow or spread.

PRODUCT FEATURES

- Broad-spectrum control of *Pythium* and *Phytophthora* diseases.
- Systemic activity provides root-to-leaf protection.
- Provides preventive and curative disease control.
- Can be used on cool- and warm-season grasses.
- 2X the concentration of competitive standard formulations.
- Excellent tank-mix partner.

LABELED USES

- Turfgrass
- Ornamental plants
- Ornamental bulb, corm, and tuber plants
- Christmas trees and Christmas tree plantations
- Non-bearing citrus trees and fruit and nut plants
- Retail nurseries
- Residential and commercial landscapes
- Interior plantscape ornamentals

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

45.3% Metalaxyl-M
(mefenoxam)

FRAC GROUP

4

FORMULATION

Microemulsion concentrate

CHEMICAL FAMILY

Acylamine

COMPARE TO

Subdue Maxx®

MODE OF ACTION

Selective inhibition of ribosomal RNA synthesis by affecting the activity of the RNA polymerases.

RESTRICTED USE

No

SIGNAL WORD

Warning

PACKAGING

4 x 0.5 gal

EPA REG NUMBER

42750-392-89442

OXEM™ 4 ME SELECT

SYSTEMIC, PREVENTIVE AND CURATIVE FUNGICIDE FOR TURF AND ORNAMENTALS.

APPLICATION INFORMATION

Label use instructions for turf, flowers and woody ornamentals	
APPLICATION AREA	LABELED DISEASES
TURF: including but not limited to golf courses, lawns, landscape areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, and athletic fields.	Downy mildew, <i>Pythium</i> blight, <i>Pythium</i> damping-off
BEDDING PLANTS: <i>Ageratum</i> , Algerian ivy, <i>Artemisia</i> , <i>Aster</i> , begonia, <i>Caladium</i> , carnation, chrysanthemum, <i>Coleus</i> , daisy, English ivy, foxglove, <i>Gaillardia</i> , geranium, <i>Impatiens</i> , marigold, pansy, petunia, <i>Phlox</i> , pinks, primrose, prostrate rosemary, <i>Salvia</i> , snapdragon, <i>Verbena</i> , <i>Vinca</i> , zinnia, including all cultivars, varieties, and hybrids of these.	Damping-off, root diseases, and stem diseases caused by <i>Pythium</i> and <i>Phytophthora</i> spp. Foliar diseases such as downy mildew and foliar diseases caused by <i>Phytophthora</i> spp., including <i>P. ramorum</i> . Downy mildew, <i>Phytophthora</i> spp., <i>Pythium</i> spp.
FLOWERS: (including bulb, corm, and tuber plants): African violet, <i>Anthurium</i> , baby's breath, carnation, chrysanthemum, columbine, <i>Delphinium</i> , Easter lily, geranium, <i>Gloxinia</i> , poinsettia, rose, including all cultivars, varieties, and hybrids.	
FOLIAGE PLANTS: <i>Aglaonema</i> , <i>Aphelandra</i> , <i>Dieffenbachia</i> , <i>Peperomia</i> , philodendron, pothos, <i>Schefflera</i> , <i>Sedum</i> , <i>Sempervivum</i> , <i>Zygocactus</i> , including all cultivars, varieties, and hybrids of these.	
WOODY ORNAMENTALS: <i>Aucuba japonica</i> , arborvitae, azaleas, boxwood, <i>Ceanothus</i> , <i>Cotoneaster</i> , dogwood, <i>Ficus</i> , "Halls" honeysuckle, <i>Ilex</i> , <i>Juniperus</i> spp., <i>Photinia</i> , <i>Pieris japonica</i> , <i>Pinus</i> spp., <i>Pittosporum</i> , <i>Rhododendron</i> , white cedar, white pine, yew, including all cultivars, varieties, and hybrids.	<i>Pythium</i> and <i>Phytophthora</i> root and crown rot. Downy mildew, <i>Phytophthora</i> spp., <i>Pythium</i> spp.
Conifers in nurseries and forest nurseries (including Christmas trees.)	<i>Phytophthora</i> root and stem diseases, <i>Phytophthora</i> foliar diseases
Conifers in plantations (including Christmas trees.)	<i>Phytophthora</i> root, stem, and foliar diseases. Foliar diseases caused by <i>Phytophthora</i> spp., including <i>P. ramorum</i>
CITRUS in nurseries and landscape plantings (non-bearing.)	Citrus foot rot, root rot, trunk canker caused by <i>Phytophthora</i> spp.
Fruit and nut plants in nurseries and landscape plantings (non-bearing.)	<i>Phytophthora</i> root, crown, and collar rot, <i>Pythium</i> root rot.



◀ See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Oxem is a trademark of Albaugh, LLC. Subdue Maxx is a registered trademark of Syngenta. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.