

HERBICIDE

TRIAD QC SELECT™

**BROAD-SPECTRUM CONTROL OF GRASS
AND BROADLEAF WEEDS.**



Triad QC Select™ combines the power of three versatile active ingredients to deliver effective postemergence weed control. Effective on both grassy and broadleaf weeds, **Triad QC Select** is the perfect option for spring break through and late season clean-up applications.

PRODUCT FEATURES

- Combines three active ingredients to help eliminate hard-to-control weeds.
- Perfect late-season clean-up product for areas that didn't receive a preemergence herbicide application in the spring.
- Can be applied as a spot application or broadcast spray.
- Controls newly germinated 1- to 2-leaf crabgrass to 1-tiller crabgrass, and when crabgrass has matured to 5-tillers or greater.
- The addition of Quinclorac delivers grass weed control through root and shoot absorption.

LABELED USES

- Residential and non-residential turfgrass
- Athletic fields
- Parks
- Golf course turf (not for use on greens, tees and collars)
- Sod farms

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

13.29% 2,4-D
9.79% quinclorac
1.67% dicamba

HRAC GROUP

2,4-D - Group 4
Quinclorac - Group 4
Dicamba - Group 4

FORMULATION

Liquid concentrate

CHEMICAL FAMILY

2,4-D - Phenoxy-carboxylic acids
Quinclorac - Quinoline carboxylic acids
Dicamba - Benzoic acids

COMPARE TO

Quincept®

MODE OF ACTION

Auxin mimic

RESTRICTED USE

No

SIGNAL WORD

Warning

PACKAGING

12 x 1 qt, 2 x 2.5 gal

EPA REG NUMBER

89442-45

TRIAD QC SELECT™

BROAD-SPECTRUM CONTROL OF GRASS AND BROADLEAF WEEDS.

APPLICATION INFORMATION

Ornamental turf and sod

Broadcast treatment: Apply 7 to 8 pints/A (112 to 128 fl oz/A) in a spray volume of 20 to 300 gal/A (0.5 to 6.9 gal/1,000 sq ft). **DO NOT** exceed 2 broadcast applications per year, excluding spot treatments. For sod, the minimum retreatment interval is 21 days.

Spot Treatment: Apply 2.6 to 2.9 fl oz per 1,000 sq ft in a spray volume of 1 gal. **DO NOT** apply greater than 16 pints of this product per acre per year. The minimum retreatment interval is 21 days.

Non-turf areas

Control of annual and perennial plants

Broadcast treatment: Apply 7 to 8 pints/A (112 to 128 fl oz/A) in a spray volume of 20 to 300 gal/A (0.5 to 6.9 gal/1,000 sq ft). **DO NOT** exceed 2 broadcast applications per year, excluding spot treatments, with a minimum retreatment interval of 30 days.

Spot treatment: Apply 2.6 to 2.9 fl oz per 1,000 sq ft in a spray volume of 1 gal. **DO NOT** apply greater than 16 pints of this product per acre per year.

Restrictions

- **DO NOT** use on golf course greens, tees, and collars.
- **DO NOT** use on lawns with desirable clovers, legumes, or on ornamentals.
- Not for use on sod farms in Arizona. For use in New York by spot treatment only (spray individual weeds only, adjust the sprayer to coarse spray to minimize wind drift, apply to the center of the weeds, and spray to lightly cover).

LABELED WEEDS

Barnyard grass	Crabgrass, large and smooth	Ground ivy	Russian pigweed	White clover
Black medic	Dandelion	Hedge bindweed	St. Johnswort	Wild carrot
Boxelder	Dock	Henbit	Shepherds purse	Wild vetch
Broadleaf dock	Dogbane	Horsetail	Smartweed	Willow
Burweed	Elderberry	Knotweed	Spurge	Woodsorrel
Carpetweed	Fall dandelion	Morning glory	Thistle	Yarrow
Chickweed	Fiddleneck	Nettle	Trailing crown vetch	Yellow rocket
Clover	Field bindweed	Oxalis	Virginia buttonweed	
		Plantain	Virginia creeper	



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Triad QC Select is a trademark of Albaugh, LLC. Quincept is a registered trademark of Nu-Farm Americas Inc. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.