

# KOVERALL<sup>®</sup>

**BROAD-SPECTRUM, CONTACT FUNGICIDE WITH ACTIVITY AGAINST A WIDE RANGE OF FUNGAL PATHOGENS.**



Koverall<sup>®</sup> is a broad-spectrum fungicide labeled for use on outdoor and greenhouse grown crops, turf, and ornamentals. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of an agricultural surfactant will enhance fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and enhanced fungicide retention during periods of wet weather. Under low disease conditions, minimum label rates per application can be used while maximum label rates and the minimum interval may be used for severe or threatening disease conditions.

## PRODUCT FEATURES

- Unique mode of action making it an excellent disease resistance management tool.
- May be applied in a foliar, seed, or chemigation application in a variety of crops.
- Effective in protecting against a wide range of fungal pathogens, including leaf spots, rusts, scabs, early blight, late blight, walnut blight, and downy mildew, in a range of crops.

### ACTIVE INGREDIENT

75% mancozeb

### FRAC GROUP

M3

### FORMULATION

Water dispersible granule

### REI

24 hours

### COMPARE TO

Manzate<sup>®</sup>

### MODE OF ACTION

Multi-site protectant

### SIGNAL WORD

Caution

### RESTRICTED USE

No

### PACKAGING

6 lb bag, 30 lb bag

### EPA REG NUMBER

42750-389 | AD021722

See label for complete application guidelines and restrictions.



# KOVERALL<sup>®</sup>

## FREQUENTLY ASKED QUESTIONS

### How should I apply Koverall on potatoes for best results?

- Apply Koverall at 1.0-2.0 lb/A for control of early blight and late blight in potatoes.
- Begin applications when plants are 4-6 inches high by applying 0.5 to 1.0 lb./acre. As the vines increase in size, apply 1.5 to 2.0 lb./acre at 5- to 10-day intervals or apply 0.75 to 1.0 lb./acre at 3- to 5- day intervals. It is directed that this product be used within an Integrated Pest Management Program. Vine-kill must occur 14 days before harvest.
- Minimum retreatment interval is 5 days.
- *Refer to label for usage restrictions.*

### How should I apply Koverall on walnuts for best results?

- Apply Koverall at 2.4 lb/A by air or ground as a tank mix for control of walnut blight.
- This product must be tank mixed with a fixed copper product that is registered on walnuts. Begin application at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage, or as needed if frequent rainfall occurs.
- Apply by ground in a minimum of 100-gal water/acre or by air in a minimum of 10 gal water/acre.
- *Refer to label for usage restrictions.*

### How should I apply Koverall on apples for best results?

- Apply 6.0 lb/A\* pre-bloom or bloom, at 1/4 to 1/2-inch greentip and continue on a 7-10 day schedule through bloom for control of *Fabraea* leaf spot, rusts and scab.
- Refer to label for extended application schedule for use in tank mixtures with systemic fungicides.
- *Refer to label for usage restrictions.*

\*Maximum per acre use rate based on thorough coverage dilute sprays.



◀ See label for complete application guidelines and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks of Albaugh, LLC. Koverall is a registered trademark of Indofil Industries. Manzate is a registered trademark of United Phosphorus Limited. Always read and follow label restrictions, precautions, and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.