



LANCER® 2EC

**BROAD-SPECTRUM INSECTICIDE, EFFECTIVE AGAINST
A RANGE OF CHEWING, FLYING AND SUCKING PESTS.**

Lancer® 2EC is a broad-spectrum synthetic pyrethroid insecticide that is effective against chewing, flying and sucking pests in a variety of crop and non-crop applications. Rate of application varies according to pest pressure, timing of sprays and field scouting. Use lower rates under light-to-moderate infestations and higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

PRODUCT FEATURES

- May be mixed with other EPA approved insecticides to improve spectrum of control in registered crops.
- Contact and residual solution that works by absorption through the insect cuticle or by ingestion for rapid knockdown.
- May be conveniently applied by ground, aerial or chemigation application systems.

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

2.0 lbs/gal bifenthrin

IRAC GROUP

3A

FORMULATION

Emulsifiable concentrate

REI

12 hrs

COMPARE TO

Capture® 2EC

MODE OF ACTION

Sodium channel modulator

SIGNAL WORD

Warning

RESTRICTED USE

Yes

PACKAGING

4 x 1 gal, 2 x 2.5 gal

EPA REG NUMBER

42750-368 | AD022824

LANCER[®] 2EC

FREQUENTLY ASKED QUESTIONS

How should I use Lancer 2EC on field corn (grain/silage), popcorn or field corn grown for seed for best (at plant use) results?

- For corn rootworm species, apply 0.3 fl oz/1,000 row feet. For cutworms, grubs, wireworms, armyworms, seed corn beetles or seed corn maggots, apply 0.15-0.3 fl oz/1,000 row feet.
- Apply as a 5-to-7-inch T-band treatment over the open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel, and centered over the row. Apply in a minimum of three gallons of finished spray per acre.
- Follow the instructions on label for mixing Lancer 2EC with water or liquid fertilizer for at-plant application.
- When applying in combination with liquid fertilizer, a jar test should be performed to ensure compatibility.

How should I adjust the at-plant application rates in corn based on row spacing?

	40" rows	38" rows	36" rows	30" rows
Lancer 2EC oz/A	3.9	4.1	4.4	5.12

Can Lancer 2EC be used as a foliar treatment on corn?

- Yes. Apply 2.1-6.4 fl oz/A, depending on insects present, in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment.
- When applying by air, use five gallons of finished spray per acre particularly when initial insect populations are heavier than normal. Additionally, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.
- For ear-attacking pests, apply Lancer 2EC just before silking and repeat as necessary to maintain control.
- Make applications for corn borer control with initial application at or shortly before egg hatch.
- For mite control, refer to specific species and state information on label. For other insect pests, apply when pests first appear and repeat as necessary.
- Do not apply within 30 days of harvest.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are actively visiting the treatment area. Protect pollinating insects by following label directions to minimize drift and to reduce risk to those organisms.



◀ See label for complete application guidelines and restrictions.

©2026 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Lancer are registered trademarks of Albaugh, LLC. Capture is a registered trademark of FMC Corporation. Lancer and Capture are restricted use pesticides. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.