



COVER XL[®]

**BROAD-SPECTRUM, PREVENTATIVE FUNGICIDE WITH
SYSTEMIC AND CURATIVE PROPERTIES RECOMMENDED
FOR THE CONTROL OF A VARIETY OF PLANT DISEASES.**



Cover XL[®] is a broad-spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. Cover XL may improve the yield and/or quality of the crop. These additional benefits are due to positive effects on plant physiology. The effects may vary according to other factors such as the crop, crop hybrid or environment. Cover XL may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products, according to label directions.

PRODUCT FEATURES

- Translocates through the plant, even through new growth, to provide preventative and curative benefits.
- Helps plants better withstand stressful growing conditions by helping build a stronger, deeper root system.
- Applications may be made by ground, air, or through chemigation providing long-lasting residual control of a variety of diseases.
- May be applied for early-season disease control and beneficial physiological effects in certain crops.

ACTIVE INGREDIENTS

1.18 lbs/gal azoxystrobin +
1.02 lbs/gal propiconazole

FRAC GROUPS

11 & 3

FORMULATION

Suspoemulsion

REI

12 hours

COMPARE TO

Quilt Xcel[®]

MODES OF ACTION

Qol inhibitor &
demethylation inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

2 x 2.5 gal, 265 gal

EPA REG NUMBER

42750-289 | AD111015

See label for complete application guidelines and restrictions.

COVER XL®

FREQUENTLY ASKED QUESTIONS

Do I need to use an adjuvant with Cover XL?

- For some labeled uses, a spreading/penetrating type adjuvant, such as non-ionic surfactant (NIS), crop oil concentrate (COC) or a blend may be added at manufacturer-recommended rates. Adjuvants that contain some form of silicone can contribute to phytotoxicity.

When should I apply Cover XL on field corn?

- From V4-V8, apply 10.5 oz/A as an early application as disease control and plant performance benefits.
- For later-season applications, apply 10.5-14.0 fl oz/A for gray leaf spot, rusts, anthracnose, and eye spot when disease first appears. If conditions remain favorable for disease, continue to apply on a 14-day schedule. For leaf blights, apply 10.5-14.0 fl oz/A when disease first appears and continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. If disease pressure is heavy, or if conditions are favorable for disease, use the higher rate.
- Do not use adjuvants or other additives after the V8 growth stage and prior to VT as use during these developmental stages may impose stress on the plant that could inhibit proper kernel development.

When should I apply Cover XL on soybeans?

- For all foliar diseases except rust, apply 14.0-21.0 fl oz/A at R3 (early pod set) and 14-21 days later at R5 (pod fill). Cover XL may be applied earlier if conditions are favorable for disease.
- For soybean rust, apply 14.0-21.0 fl oz/A at first indication that disease is in the area. Preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter intervals when diseases are present in the field and incidence is less than 2%. If incidence is greater than 2% or if the disease is mid-canopy, control will not be acceptable. Use proper scouting procedures to maximize effectiveness of the application.
- On certain varieties, Cover XL applications may cause crinkled, smaller and/or greener leaves. Yield has not been reduced due to Cover XL applications.
- Apply up to Stage R6.



◀ See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Cover XL are registered trademarks of Albaugh, LLC. Quilt Xcel is a registered trademark of a Syngenta Group Company. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.