

ACCEPTED  
FOR REGISTRATION  
Jan 14, 2026

# DIAMID™ T&O

## FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL

**FOR USE IN KINGS, QUEENS, NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE  
ONLY ON GOLF COURSE TURF**

DOC ID# 603298

**Diamid™ T&O  
For Control of White Grubs, Turf Caterpillars and Billbugs on  
Golf Course Turfgrass Only**

**EPA Reg. No. 89442-70  
EPA SLN No. NY-250003**

**This label expires and must not be distributed or used in accordance with  
this SLN registration after December 31, 2029**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

The following restrictions are required to permit use of Diamid™ T&O in the State of New York:

- Product is restricted use by certified applicators only in New York State
- **Golf course greens and tee boxes:** DO NOT apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- **Golf course fairways:** DO NOT apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch),
- **Other use sites:** DO NOT apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

### **APPLICATION RATES FOR GOLF COURSES**

For maximum residual effectiveness or for optimal control of large pest infestations, apply Diamid™ T&O at up to 10 fluid ounces per acre (= 0.23 fluid ounces per 1,000 square feet or 0.26 lb AI/A) to control the pests listed in Table 1 below. Apply Diamid™ T&O in sufficient water to provide optimal distribution in the treated area. Use properly calibrated application equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift.

**Restriction:** Do not apply more than 10.0 fluid ounces (equivalent to 0.26 lb of active ingredient) of product per acre per year in broadcast applications to golf turfgrass.

## APPLICATION TIMING

**Billbugs:** Apply Diamid™ T&O when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrass. The higher rate listed in Table 1 may be required for applications made after early May or when billbug species other than bluegrass billbug are present. An application at this time will also provide white grub control.

**White Grubs:** Apply Diamid™ T&O from early April to early September for preventative and early curative control of all the major white grub species infesting turfgrass. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. The higher rate listed for white grub control in Table 1 may be required [for early season (April and May) applications where a long residual is required or] in late August or early September when less sensitive mid-instar grubs are present at the time of application. Irrigate turf immediately after application or allow rainfall to move the product into the soil.

**Turf Caterpillars:** Apply Diamid™ T&O preventatively or after caterpillars have been observed. To ensure best control, delay water (irrigation) or mowing for 24 hours after application.. If the area being treated is maintained at a mowing height of greater than one inch, then the higher rate listed in Table 1 may be required during periods of high pest pressure.

**TABLE 1: Turf Application Rates**

Target Pest	Diamid™ T&O Turf Application Rates		
	Product (fl oz) per Acre	Product (fl oz) per 1,000 sq ft	Ib AI/A
Billbugs	4 - 10	0.09 – 0.23	0.104 - .026
Turf Caterpillars (including armyworms, cutworms and sod webworms)	1 - 2	0.025 - 0.045	0.026 - 0.052
White Grubs (including Aphodius spp., Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, May/June beetles ( <i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle and southern masked chafer)	4 - 8	0.09 to 0.185	0.104 - 0.208

24(c) Registrant:

Prime Source, a division of Albaugh, LLC  
1525 NE 36th Street  
Ankeny, IA 50021

SLN\_NY\_010625