

MITICIDE

# HEXAMITE<sup>®</sup> 1AQ SELECT

MITICIDE FOR USE ON TURF, ORNAMENTALS,  
AND NON-BEARING FRUIT TREES AND VINES.



Hexamite<sup>®</sup> 1AQ Select controls mites through activity on eggs and immature stages through either direct contact with the spray or from contact with treated plant surfaces. Ovicidal activity controls newly deposited mite eggs and eggs that are laid after application. Hexamite 1AQ Select is highly effective in controlling immature motile stages of target mite species that have been sprayed or have moved onto treated surfaces. Adult mites are not directly affected. The degree and duration of control is dependent on the use rate, mite species, growth stage, and climatic conditions at the time of application.

## PRODUCT FEATURES

- Long-lasting miticide with residual control for up to two months.
- Unique mode of action provides effective activity on eggs and early stages of development.
- Multiple use sites.

## LABELED USES

- Turfgrass
- Ornamentals
- In and around residential, institutional, public, commercial, agricultural, and industrial buildings.

See label for complete application guidelines and restrictions.

### ACTIVE INGREDIENT

11.8% hexythiazox

### IRAC GROUP

10A

### FORMULATION

Aqueous suspension

### CHEMICAL FAMILY

Thiazolidinone

### COMPARE TO

Hexygon<sup>®</sup> IQ

### MODE OF ACTION

Controls mite growth through activity on eggs and early stages of development.

### RESTRICTED USE

No

### SIGNAL WORD

Caution

### PACKAGING

12 x 1 qt

### EPA REG NUMBER

89442-59

# HEXAMITE® 1AQ SELECT

MITICIDE FOR USE ON TURF, ORNAMENTALS, AND NON-BEARING FRUIT TREES AND VINES.

## LABELED MITES

Arborvitae spider mite	Clover mite	Pecan leaf scorch mite	Tetranychid mites
Banks grass mite	European red mite	Southern red mite	Two-spotted spider mite
Brown wheat mite	Honeylocust spider mite	Spruce spider mite	Willamette mite
Citrus red mite	Pacific spider mite	Strawberry spider mite	

## APPLICATION DIRECTIONS FOR ORNAMENTALS AND TURF

CROP	PEST	RATE	USE DIRECTIONS
ORNAMENTAL PLANTS AND NON-BEARING FRUIT TREES AND VINES Growing in nurseries, greenhouses, shadehouses and Christmas tree plantations	Arborvitae spider mite Banks Grass mite Brown wheat mite Clover mite European red mite Honeylocust spider mite Pacific spider mite Southern red mite Spruce spider mite Strawberry spider mite Two-spotted spider mite Willamette mite	12 - 24 oz/A OR	When using high-volume dilute sprays, do not exceed the maximum per acre rate or apply less than the minimum labeled rate. For concentrate equipment, use the recommended rate per acre. Apply to lawns and turf at the first signs of egg deposition.
ESTABLISHED ORNAMENTAL LANDSCAPE PLANTINGS Including those in: Interiorscapes, residences, public areas, commercial areas, institutional areas, industrial areas, rights of way and other easements, and recreational sites such as campgrounds, golfcourses, parks, and athletic fields		4 - 8 oz/100 gal OR	
		0.28 - 0.55 oz/ 1000 sq ft	
ORNAMENTAL LAWNS AND TURF (including residential) except California			
NON-RESIDENTIAL TURFGRASS (including turf grown for sod, turf grown to produce seed, or turf for research purposes) Except California			



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh, the A logo, and Hexamite are registered trademarks of Albaugh, LLC. Hexygon is a registered trademark of Gowan Company. Always read and follow label restrictions, precautions, and directions for use. Products may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.