

FIFRA 2(ee)



Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021

MALIBU™ PLUS D

EPA Reg. No. 83100-78

2(ee) Recommendation

Control of Susceptible Weeds Pre or Post Planting of Certain Annual and Perennial Grasses

ATTENTION

This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for MALIBU™ PLUS D herbicide.

Directions for Use

This 2(ee) recommendation is limited to the use sites listed on the main label for MALIBU™ PLUS D specialty herbicide. Refer to the main label for use sites.

MALIBU™ PLUS D controls susceptible broadleaf plants both pre and postemergence. See the MALIBU™ PLUS D main label for a list of susceptible weeds.

PRECAUTIONS

Postemergence applications on newly emerged or established sorghum-sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with MALIBU™ PLUS D applications.

Pre-plant applications for control of susceptible emerged weeds

Apply MALIBU™ PLUS D at a maximum of 1.5 pts/acre with a non-ionic surfactant to control susceptible weeds no sooner than 15 days prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass. MALIBU™ PLUS D can also be used at 1.5 pts/acre prior to planting sorghum, sudangrass, teff, crabgrass, and pearl millet, but a minimum 30-day interval must be observed between the application and planting.

Post planting applications for control of susceptible emerged weeds

Apply MALIBU™ PLUS D at a maximum of 1.5 pts/acre with a non-ionic surfactant to tall fescue, orchardgrass, timothy, and annual ryegrass after seedling plants have at least reached the V3 stage (three collared leaves).