

CLOMATE™ 3ME

**MICROENCAPSULATED FORMULATION WITH
VERSATILE APPLICATION FOR CONTROL OF ANNUAL
GRASSY WEEDS IN RICE AND OTHER CROPS.**



Apply Clomate™ 3ME alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. Use nozzles suitable for broadcast boom or banded application of herbicides, observing specific label instructions to reduce spray drift. Tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers need to be no finer than 50-mesh. Clomate 3ME may be used as a preemergence soil surface applied treatment from 14 days prior to planting or up to 7 days after planting, but prior to weed emergence. This product may be applied after planting as an early postemergence treatment to rice through the 5-leaf stage.

Clomate 3ME is not approved for use in California.

PRODUCT FEATURES

- Microencapsulated formulation minimizes movement away from the site of application.
- Effective foundation herbicide for control of a range of grassy weeds in rice, including barnyardgrass, broadleaf signalgrass, panicum, and sprangletop.
- Excellent fit in an overall weed management program in Clearfield® rice systems to reduce competition from early season weeds.
- Flexible timing allows for pre-plant, at-plant or early postemergence applications up to five-leaf rice.

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

3.0 lbs/gal clomazone

HRAC GROUP

13

FORMULATION

Microencapsulated

REI

12 hours, refer to label for exceptions

COMPARE TO

Command® 3ME

MODE OF ACTION

Carotenoid biosynthesis inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

2 x 2.5 gal

EPA REG NUMBER

45002-60 | AD092223

CLOMATE™ 3ME

FREQUENTLY ASKED QUESTIONS

How should I apply Clomate 3ME in rice for best results?

Preemergence surface broadcast applications

- Apply Clomate 3ME as a preemergence surface broadcast application 14 days prior to planting or up to 7 days after planting, but prior to weed emergence, using ground equipment in a minimum of 10 to 40 gallons of water per acre at the rate of 10.7 to 34.1 fl. oz. product (0.25 to 0.8 lb. active) per acre depending upon the soil texture. Refer to the label for specific rates and weeds controlled. For heavy soils use the higher specified rate, otherwise less than desirable weed control may result.

Early postemergence applications

- Apply Clomate 3ME after planting as an early postemergence treatment to rice through the five-leaf stage to provide preemergence and residual control of grass weeds. Use ground equipment with nozzles that produce a coarse spray and a minimum of 10-40 gallons of water per acre. Refer to table below for specific rates and weeds controlled. For control of existing grass present at the time of application include a postemergence herbicide registered for the control of grass species in rice. Consult postemergence herbicide label for specific directions regarding use rates and stage of weeds and crop.
- Partial weed control may result if levees are pulled after this product has been applied. Additional use of labeled postemergence herbicide applications may be required.
- Refer to label for split-application instructions, rotational crop restrictions, and usage restrictions.
- Clomate 3ME is not approved for use in California or in water-seeded rice.

At what rate should I apply Clomate 3ME on rice?

Soil Texture	Broadcast Rates per Acre
Coarse (light) soils: sand, loamy sand, sandy loam	10.7-14.1 fl. oz. product (0.25-0.33 lb ai)
Medium soils: loam, silt, silt loam, sandy clay, sandy clay loam	17.1-21.3 fl. oz. product (0.4-0.5 lb ai)
Fine (heavy) soils: silty clay, clay loam, silty clay loam, clay	21.3-34.1 fl. oz. product (0.5-0.8 lb ai)

Select lower to higher rates based on lighter to heavier soils.

Can I replant rice in fields treated with Clomate 3ME?

- Yes. If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with Clomate 3ME. When tank mixing with a labeled product, refer to the replant instructions for that product. When a tank mix is used, refer to the product's label for any additional rotational crop instructions.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are trademarks of Albaugh, LLC. Command is a registered trademark of FMC Corporation. Clearfield is a registered trademark of BASF Corporation. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status. All products are EPA registered.