

NON-SELECTIVE HERBICIDE

THUNDER MASTER[®]

ROUNDUP READY^{*} SOYBEANS AND ALFALFA, OR AS A FALL OR SPRING WEED BURNDOWN IN CLEARFIELD^{*} CORN.



Thunder Master[®] provides burndown and residual control of many broadleaf and grass weeds in soybeans and Roundup Ready[®] alfalfa. May be applied postemergence to Roundup Ready soybeans and Roundup Ready alfalfa for seasonlong weed control, or as a fall or spring burndown in CLEARFIELD[®] corn. Thunder Master is effective for difficult-to-control weeds common to no-till production, including marestail. When applied postemergence, Thunder Master not only controls existing broadleaf and grass weeds, it also provides control of susceptible weeds that may emerge after application.

PRODUCT FEATURES

- Burndown plus the residual for control of many broadleaf and grass weeds in Roundup Ready soybeans and alfalfa.
- Two modes of action to manage herbicide resistance.
- Flexibility of use, may be applied preplant burndown, and early postemergence.
- Apply postemergence in Roundup Ready soybeans and Roundup Ready alfalfa for season-long control.
- Excellent residual control, resulting in fewer trips across the field.

ACTIVE INGREDIENTS

2.7 lbs/gal imazethapyr +2.0 lbs/gal glyphosate

HRAC GROUPS

FORMULATION Soluble concentrate

REI 48 hours

COMPARE TO Extreme®

MODES OF ACTION ALS inhibitor, EPSP synthase inhibitor

SIGNAL WORD

PACKAGING 2 x 2.5 gal, 30 gal, NR 265, Bulk

EPA REG NUMBER 42750-147 | AD060716



See label for complete application guidelines and restrictions.



FREQUENTLY ASKED QUESTIONS

How should I apply Thunder Master herbicide for best results?

Postemergence

- A non-ionic surfactant and spray grade ammonium sulfate fertilizer must be added to the spray solution for optimum weed control activity. *Refer to MIXING INSTRUCTIONS on page 6-7 of the product label for specific instructions.*
- For maximum weed control, cultivate 7-10 days following a postemergence application of Thunder Master. This timely cultivation will enhance residual weed control, especially under dry conditions.
- Apply to actively growing weeds.
- Apply a minimum of one hour before rainfall or overhead irrigation.
- Use only flat-fan nozzles for postemergence applications.
- Delay application for 48 hours from the time the temperature increases above 50F, if air temperature has been below 50 F for 10 hours or more to improve weed control and reduce crop response.

Spraying Instructions

- Ground applications should be made in 10 or more gallons of water per acre with a recommended spray pressure of 20-40 psi. Use a minimum of 20 gallons of water per acre when applying Thunder Master to minimum or no-till crops. Use a higher gallonage for fields with dense vegetation or heavy crop residue.
- If applying with a low-volume sprayer, apply a minimum of 10 gallons per acre with nozzle pressure between 40-60 psi for optimum coverage.
- Aerial applications should be made in five or more gallons of water per acre. When applied postemergence, the addition of non-ionic surfactant AND fertilizer solution are required for optimum weed control.
- Refer to label for specific application information by crop.

With the resistance that is so prevalent with this class of chemistry, what does Thunder Master bring to the table?

• Residual grass control, as well as control of regionally challenging broadleaf weeds using two active ingredients.



See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo, Thunder Master and Extreme are registered trademark of Albaugh, LLC. CLEARFIELD is a registered trademark of BASF Corporation. Roundup Ready is a registered trademark of Monsanto Technology LLC. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.

