

THUNDER

PROVIDES LONG-LASTING CONTROL OF ANNUAL GRASS AND BOTH LARGE- AND SMALL-SEEDED BROADLEAF WEEDS WITH APPLICATION FLEXIBILITY.



Thunder* herbicide kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum activity. When adequate soil moisture is present, Thunder will provide residual control of susceptible germinating weeds. Activity on established weeds will depend on the weed species and the location of its root system in the soil. Occasionally, internode shortening and/or temporary yellowing of the crop may occur following Thunder applications. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1-2 weeks.

PRODUCT FEATURES

- Proven control of tough broadleaf and grass weeds.
- Broad crop label that includes alfalfa, clover, peas and beans, field corn*, peanuts and soybeans.
- May be applied preemergence (surface application), postemergence, soil applied with liquid fertilizers, in no-till or reduced tillage or preplant incorporation applications.
- *Apply Thunder herbicide only on field corn hybrids (Clearfield® corn) warranted by the seed company to possess resistance/tolerance to direct application of Thunder.

ACTIVE INGREDIENT

2.0 lb/gal imazethapyr

HRAC GROUP

2

FORMULATION

Soluble concentrate

REI

4 hours

COMPARE TO

Pursuit*

MODE OF ACTION

ALS inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

4 x 1 gal

EPA REG NUMBER

42750-146 | AD070617

See label for complete application guidelines and restrictions.





FREQUENTLY ASKED QUESTIONS

How should I apply Thunder herbicide for best results?

Postemergence

- An adjuvant (either crop oil concentrate or a surfactant) and a
 nitrogen-based fertilizer must be added to the spray solution
 for optimum weed control activity. Refer to the ADJUVANTS
 section under MIXING INSTRUCTIONS on page 7 of the
 product label for specific instructions. NOTE: Mixing Thunder
 with a fertilizer solution is prohibited in California.
- For maximum weed control, cultivate 7-10 days following a
 postemergence application of Thunder. This timely cultivation
 will enhance residual weed control, especially under
 dry conditions.
- Apply to actively growing weeds, before they exceed a height of three inches, unless otherwise indicated. Delay application until the majority of the weeds are at the recommended growth stage.
- Apply a minimum of one hour before rainfall or overhead irrigation.
- Use only flat-fan nozzles for postemergence applications.
- Delay application for 48 hours from the time the temperature increases above 50F, if air temperature has been below 50F for 10 hours or more to improve weed control and reduce crop response.
- Refer to label pages 8-12 for specific application information.

Soil application

- Adequate soil moisture is required for optimum activity and to move Thunder into the germination zone. Amount of moisture required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2" is normally adequate.
- If adequate moisture is not received within seven days after treatment, a cultivation is recommended to control escaped weeds.
- Refer to label pages 8-12 for specific application information.



 \blacktriangleleft See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Thunder are registered trademarks of Albaugh, LLC. Clearfield and Pursuit are registered trademarks of BASF Corporation. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.

